arXiv:1406.0385v1 [math.NT] 2 Jun 2014

DENSITY OF SELF-DUAL AUTOMORPHIC REPRESENTATIONS OF $GL_N(\mathbb{A}_{\mathbb{Q}})$

A Dissertation

Submitted to the Faculty

of

Purdue University

by

Vítězslav Kala

In Partial Fulfillment of the

Requirements for the Degree

of

Doctor of Philosophy

August 2014

Purdue University

West Lafayette, Indiana

ACKNOWLEDGMENTS

I want to greatly thank my advisor Professor Shahidi for all his valuable advice, support and encouragements during my work with him. I wish to thank Professors Goldberg, Liu, and McReynolds for agreeing to serve on my committee and for their help and useful suggestions. I am grateful to Professors Arthur, Jiang, Rohrlich, and J.-K. Yu, and to Britain Cox, Radhika Ganapathy, Yeansu Kim, Jing Feng Lau, Manish Mishra, Abhishek Parab, and Sohei Yasuda, for interesting mathematical discussions and suggestions which have influenced and improved this thesis. I also want to thank many other professors and staff at Purdue and Charles University for their help and support. Let me mention only Professors Lipshitz and Drápal, whose mentorship and support have been very important for me. During my first three years at Purdue I have been fully supported by the International Fulbright Science and Technology Award – I greatly appreciate this opportunity. And finally, I thank my family and all my friends, both home in the Czech Republic and in the US. Without you it wouldn't have been so much fun!

TABLE OF CONTENTS

Page

LIST OF TABLES iv

ABSTRACT v

1 Introduction 1

2 Background 9
2.1 Basic Definitions and Notation 9
2.2 Langlands Functoriality and Reciprocity 10
2.3 Weyl's Law 15
2.4 L-Functions and Irreducibility of Induced Representations 18

3 Laplacian Eigenvalues 21
3.1 Casimir Element 21
3.2 Infinitesimal Character 22
3.3 Behavior under Functoriality 23

4 Depth 27
4.1 Background 27
4.2 Bounds on Depth 29
4.3 Local Descent 33
4.4 Depth Preservation and Wildly Ramified Groups 36
4.5 Restriction and Depth 37

5 Upper Bound 40
5.1 Tools 40
5.2 Statement and Proof of Theorem 5.2.1 42

6 Lower Bound 44
6.1 Non-Cuspidal Lifts 44
6.2 Statement and Proof of Theorem 6.2.1 47

LIST OF REFERENCES 51

VITA 54

LIST OF TABLES

Table Page

2.1 Functoriality 12

2.2 Dimensions 17

ABSTRACT

Kala, Vítězslav Ph.D., Purdue University, August 2014. Density of Self-Dual Automorphic Representations of $GL_n(\mathbb{A}_{\mathbb{Q}})$. Major Professor: Freydoon Shahidi.

We study the number $N_{\mathrm{sd}}^K(\lambda)$ of self-dual cuspidal automorphic representations of $GL_N(\mathbb{A}_{\mathbb{Q}})$ which are K-spherical with respect to a fixed compact subgroup K and whose Laplacian eigenvalue is $\leq \lambda$. We prove Weak Weyl's Law for $N_{\mathrm{sd}}^K(\lambda)$ in the form that there are positive constants c_1, c_2 (depending on K) and d such that $c_1\lambda^{d/2} \leq N_{\mathrm{sd}}^K(\lambda) \leq c_2\lambda^{d/2}$ for all sufficiently large λ. When $N = 2n$ is even and K is a maximal compact subgroup at all places, we prove Weyl's Law for the number of self-dual representations, i.e., $N_{\mathrm{sd}}^K(\lambda) = c\lambda^{d/2} + o(\lambda^{d/2})$. These results are based on considering functorial descents of self-dual representations Π to quasisplit classical groups $\mathbf{G}$.

In order to relate the properties of representations under functoriality, we discuss the infinitesimal character of the real component Π_∞, which determines the Laplacian eigenvalue. To relate the existence of K-fixed vectors, we study the depth of p-adic representations, proving a weak version of depth preservation. We also consider the explicit construction of local descent, which allows us to improve the results towards depth preservation for generic representations.

1. Introduction

One of the important problems in modern number theory is the study of automorphic representations, i.e., representations which appear in the spectral decomposition of $L^2(\mathbf{G}(F)\backslash\mathbf{G}(\mathbb{A}_F))$ for a reductive group $\mathbf{G}$ over a number field F.

Among the first incarnations of this idea was the theory of modular and Maass forms, which are essentially just automorphic forms on $GL_2(\mathbb{A}_{\mathbb{Q}})$. It is not hard to construct examples of modular forms, but Maass forms are much harder to handle concretely – in fact, at first it was not clear if they existed at all. Their existence and asymptotics were first proved by Selberg [30] in the form of Weyl's Law, which he deduced from his trace formula.

In the classical setting, Weyl's Law provides the asymptotic estimates for the number of eigenvalues of the Laplacian operator on a compact Riemannian manifold. Automorphic forms are, among other things, eigenfunctions of the Laplacian on locally symmetric spaces $A_{\mathbb{R}}G(F)\backslash G(\mathbb{A}_F)/K$ for suitable compact subgroups K of $G(\mathbb{A}_F)$. Weyl's Law is expected to hold in some generality even in this setting, and has been proven in a number of cases, e.g., by Müller [27] for $GL_N(\mathbb{A}_{\mathbb{Q}})$ and by Lindenstrauss-Venkatesh [23] for split adjoint semisimple groups $\mathbf{G}$. See Section 2.3 for a precise statement and more background information on Weyl's Law.

The principle of functoriality, predicting transfers of automorphic representations between certain pairs of groups G_1 and G_2, plays a crucial role in the Langlands program. There has been quite a lot of progress recently in establishing functoriality from (quasisplit) classical groups to GL_N, first by Cogdell, Kim, Piatetski-Shapiro, and Shahidi [8] for generic representations and then by Arthur [3] for all representations. In these cases, the images of functorial lifts are among the self-dual automorphic representations of $GL_N(\mathbb{A}_{\mathbb{Q}})$, as it follows from the work of Ginsburg, Rallis, and Soudry [13]. See Section 2.2 for more information on functoriality.

However, being self-dual is a fairly restrictive condition. Thus we may ask how many self-dual representations are there? Do they have density zero among all representations?

In a local setting, Adler [2] gave explicit formulas for the number of depth zero self-dual supercuspidal representations of GL_N over a local field. For (global) Galois representations, Rohrlich [29] has proved that the density is zero for representations $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \to GL_N(\mathbb{C})$ when $N = 1$ or 2 and obtained some conditional results for $N = 3$.

The main result of our thesis is the following theorem, concerning the number of self-dual cuspidal automorphic representations of $GL_N(\mathbb{A}_\mathbb{Q})$:

Theorem 1.0.1 *Let $N \in \mathbb{N}$. For $M \in \mathbb{N}$ denote by $K(M)$ the principal congruence subgroup of $GL_N(\mathbb{A}_\mathbb{Q})$, i.e., $K(M) = K_\infty \times \prod K_p$ with $K_\infty = O(N)$ and K_p consisting of matrices congruent to the identity modulo $p^{m(p)}$, where $M = \prod p^{m(p)}$.*

For $\lambda > 0$, let

$$N_{\mathrm{sd}}^{K(M)}(\lambda) = \sum_{\mathrm{sd},\ \lambda(\Pi)\leq\lambda} \dim \Pi^{K(M)},$$

the sum ranging over self-dual cuspidal automorphic representations Π of $GL_N(\mathbb{A}_\mathbb{Q})$ with Laplacian eigenvalue $\leq \lambda$.

Then there are positive constants c_1 and c_2 (depending on K, but not on λ) such that, for sufficiently large λ, one has

$$c_1\lambda^{d/2} \leq N_{\mathrm{sd}}^{K(M)}(\lambda) \leq c_2\lambda^{d/2},$$

where $d = n^2 + n$ for $N = 2n + \varepsilon$ with $\varepsilon = 0, 1$.

On the other hand, by Weyl's Law (Theorem 2.3.1) the number of all cuspidal automorphic representations of $GL_N(\mathbb{A}_\mathbb{Q})$ (counted similarly) is asymptotically $c\lambda^{D/2}$ for $D = (N^2 + N - 2)/2$. Hence when $N \neq 2$, the density of self-dual cusp forms is indeed zero. When $N = 2$, self-dual cusp forms have positive density due to the fact that $SO_3 = PGL_2$ – the lifts from this group to GL_2 provide for the positive proportion of self-dual representations.

To prove Theorem 1.0.1, we prove upper and lower bounds as Theorems 5.2.1 and 6.2.1. The general idea of the proof is to consider the descent π of each self-dual cuspidal automorphic representation Π to one of the quasisplit classical groups $\mathbf{G}(\mathbb{A}_{\mathbb{Q}})$ and to use results towards Weyl's Law on $\mathbf{G}(\mathbb{A}_{\mathbb{Q}})$. For that we need to relate the relevant properties of Π and π, mainly the Laplacian eigenvalue and the existence of K-fixed vectors.

The Laplacian eigenvalue of an automorphic representation σ is determined by the infinitesimal character μ of the real component σ_∞, namely, by evaluating μ on the Casimir element Ω. For real groups we have the Langlands reciprocity and there is an explicit description of the infinitesimal character of σ_∞ in terms of the associated parameter $\varphi : W_{\mathbb{R}} \to {}^L G$ (here $W_{\mathbb{R}}$ is the Weil group and ${}^L G$ is the Langlands dual group, defined in Section 2.2). In our setting of functoriality from classical groups to GL_N, the parameter of the functorial lift is just obtained as the composition

$$\iota \circ \varphi : W_{\mathbb{R}} \to {}^L G \hookrightarrow {}^L GL_N.$$

This enables us to conclude that the infinitesimal character, and hence also the Laplacian eigenvalue, are essentially preserved under functoriality, as proved in Theorem 3.3.2.

We also need to relate the existence of K-fixed vectors for π and Π. This is a purely local question, but it is still quite involved, as we need to use the local notion of depth to obtain Proposition 5.1.1. We shall discuss this in more detail shortly, but first let's explain how we use these results to prove Theorem 1.0.1, starting with the upper bound.

Let $K = K(M)$ be a principal congruence subgroup as above and let $\lambda > 0$. We need to estimate $N_{\mathrm{sd}}^K(\lambda) = \sum_{\mathrm{sd},\ \lambda(\Pi)\leq\lambda} \dim \Pi^K$, the sum ranging over self-dual cuspidal automorphic representations Π of $GL_N(\mathbb{A}_{\mathbb{Q}})$ with Laplacian eigenvalue $\leq \lambda$. Each of these self-dual Π descends to one of the split or non-split symplectic or special orthogonal groups, determined by the poles of the symmetric square and exterior square L-functions and by the central character χ in the case of the quasisplit even

orthogonal groups. The restriction of χ to the compact subgroup K has to be trivial, and so there are only finitely many possibilities for χ and (by class field theory) thus only finitely many possibilities for the corresponding quasisplit group SO^*_{2n} (see Lemma 5.1.2).

Thus we can express $N^K_{\mathrm{sd}}(\lambda)$ as a finite sum $\sum_G N^K_{\mathrm{sd},\mathbf{G}}(\lambda)$ over different classical groups to which Π can descend. By 3.3.2 and 5.1.1, we have a corresponding bound on the Laplacians and compact subgroups for the descents π on each of these groups G, and so we can estimate $N^K_{\mathrm{sd},\mathbf{G}}(\lambda)$ from above by the number of all cuspidal automorphic representations on G as in Weyl's Law. It is not yet known on classical groups, but (after translating from the adelic to classical language) Donnelly's result [10] gives an upper bound, which is all we need.

The idea of the proof of the lower bound is similar. It suffices to consider only the case when K is a maximal compact subgroup at all places, which somewhat simplifies the situation, but we have to overcome two additional difficulties: First, a cuspidal automorphic representation on a classical group need not lift to a cuspidal representation on GL_N. However, we show that asymptotically these representations have density zero (Proposition 6.1.2). And second, we need to obtain an estimate on the size of (K-fixed elements of) global packets. In the case when K is maximal at all places, it in fact follows from Arthur's work that each global packet has at most one K-spherical element (Lemma 2.2.5). The lower bound then follows from Weak Weyl's Law for $\mathbf{G}(\mathbb{A}_{\mathbb{Q}})$ (see Theorem 2.3.2 and the discussion surrounding it).

In the proof of the lower bound, we are relying on Arthur's results [3], which are conditional on the stabilization of the twisted trace formula. To prove the upper bound, it suffices to consider only functoriality for generic representations, which is available by [9] unconditionally.

Note that when $N = 2n$ and K is a maximal compact subgroup, Weyl's Law is known for $\mathbf{G}(\mathbb{A}_{\mathbb{Q}}) = SO_{2n+1}(\mathbb{A}_{\mathbb{Q}})$ [23] and we can prove the exact asymptotics (without an error term) for $N^K_{\mathrm{sd}}(\lambda)$ (Corollary 6.2.2). It should be possible to obtain this result also when N is odd – but extending this to the case when K need not be

maximal seems to be difficult. One can try to study and estimate the twisted trace formula in a similar way as in the proof of Weyl's Law for GL_N by Lapid-Müller [22]. This is currently being pursued by Abhishek Parab under the supervision of Professor Shahidi.

Let us now turn to the existence of vectors fixed by open compact subgroups in the local setting. We need these results in the proof of a local result, Corollary 4.2.4, which then implies a global statement, Proposition 5.1.1.

Now take F to be a p-adic field, $\mathbf{G}$ a connected reductive group defined over F and π an irreducible admissible representation of $G = \mathbf{G}(F)$. In the cases when $\mathbf{G}$ is GL_N or a classical group SO_N, Sp_{2n} or SO^*_{2n}, the local Langlands correspondence has been proved by Harris–Taylor [15], Henniart [16], and Arthur [3] (in a slightly weaker form for even orthogonal groups). Thus we can associate to π a representation of the Weil-Deligne group $\varphi : W'_F \to {}^L G$.

Fix an open compact subgroup K of G. How can we describe the existence of K-fixed vectors in π in terms of properties of φ? (This would enable us to see what happens to the fixed vectors under functoriality, as in Corollary 4.2.4.)

For $\mathbf{G} = GL_N$ the answer is due to Jacquet, Piatetski-Shapiro, and Shalika [17]. They define a filtration of $GL_N(F)$ by certain "mirabolic" subgroups M_j and show that when π is generic, the smallest j such that $\pi^{M_j} \neq \{0\}$ is the analytic conductor of π given by the associated ε-factor. The L-functions and ε-factors are preserved under local Langlands correspondence, and so this provides an answer to our question.

The case of a classical group G is complicated by two facts: first of all, not every tempered representation is generic. But even more seriously, we do not have a suitable analogue of the mirabolic subgroups. A more convenient notion is that of depth, defined by Moy and Prasad [25], [26] via filtration by subgroups of G attached to points in the Bruhat-Tits building. The definition itself is somewhat technical (see Section 4.1), but the main properties of depth are that the depth of a representation π is a non-negative rational number and that depth is preserved under subquotients

of parabolic induction and the Jacquet functor. That is, if M is a Levi subgroup of G and σ an irreducible admissible representation of M, then the depth of every irreducible subquotient of the normalized parabolic induction $\mathrm{Ind}_{MN}^{G}\sigma$ is equal to the depth of σ (and similarly for the Jacquet functor).

Finally, we expect to have depth preservation: define the depth of a Weil-Deligne representation $\varphi : W_F' \to {}^L G$ as $\inf\{r \mid \varphi(I_s) = 1 \text{ for all } s > r\}$, where I_s is the filtration of the inertia subgroup I of $W_F \subset W_F'$. Then depth (π) is expected to be equal to depth (φ). J.-K. Yu [35] has proved this for GL_N and for tamely ramified tori; however, it is not true for example for the non-split torus $SO_2^*(\mathbb{Q}_p)$ corresponding to a quadratic extension $F/\mathbb{Q}_p$ which is ramified at $p = 2$ (see Corollary 4.4.2). The fact that depth need not be preserved when residual characteristic is small has already been observed by Gross-Reeder [14]; Reeder-Yu [28] recently provided more examples of this phenomenon.

We use the notion of depth to relate the existence of fixed vectors for a self-dual representation Π of $GL_N(F)$ and its descent π to a classical group G. For that, we prove Proposition 4.2.3, which provides the result needed in counting self-dual representations.

We also look at proving the depth preservation for classical groups. As we have noted before, it is not true in general for non-split even orthogonal groups, but we prove at least one inequality for generic representations in Theorem 4.3.1. The proof is based on the local descent construction of Jiang and Soudry [18]. Since depth is preserved under parabolic induction, we can assume that Π and π are supercuspidal, in which case the local descent gives an explicit way of constructing π from Π in terms of certain inductions and Jacquet quotients, which preserve depth. The obstruction to proving depth preservation by this method is that one of the steps in the construction consists of taking the restriction of a representation to a subgroup, which may decrease the depth. See Section 4.3 for the proof and details about local descent (in the case of $\mathbf{G} = SO_{2n+1}$).

The restriction that appears in the local descent to SO_{2n+1} is that of a representation of $SO_{2n+2}(F) \times (GL_1(F))^{n-1}$ to the first component $SO_{2n+2}(F)$ and then to $SO_{2n+1}(F) \subset SO_{2n+2}(F)$. In Section 4.5 we study a part of this problem, i.e., the restriction from $SO_{2n+2}(F)$ to $SO_{2n+1}(F)$ and show that depth does not decrease under it. To prove this, we use some results on irreducibility of parabolically induced representations which follow from the Langlands-Shahidi method and are stated in Section 2.4. We hope to work on this problem further in the future and hopefully obtain full depth preservation for quasisplit classical groups at least in some cases.

Our method should work quite generally for obtaining upper bounds on the number of functorial lifts from $G(\mathbb{A}_{\mathbb{Q}})$ to $GL_N(\mathbb{A}_{\mathbb{Q}})$ whenever functoriality is known at least for generic representations. This is the case for lifts from unitary and general spin groups to GL_N ([24], [4]); the image of functoriality is formed by conjugate self-dual and essentially self-dual representations, respectively. As was the case in this thesis, proving lower bounds is much more delicate and requires more detailed information about the image of functoriality and the global packets. This seems to be available at least in the case of unitary groups. We plan to undertake this research in more detail in the future. It also seems possible to extend these results to other number fields besides $\mathbb{Q}$, with the caveat that Weyl's Law is not known in most of these cases.

One may wonder how our results on the number of self-dual automorphic representations relate to Rohrlich's results [29] concerning Galois representations. An automorphic representation attached to a Galois representation will have Laplacian eigenvalue 0; essentially, Rohrlich is considering the asymptotics when the Laplacian eigenvalue (which corresponds to the weight in classical setting) is fixed (and small) and the level of the compact subgroup goes to infinity. On the other hand, our results concern the asymptotics when the level is fixed and the Laplacian eigenvalue goes to infinity. This also explains the seeming disparity between our result that

self-dual cusp forms on GL_2 have positive density, whereas 2-dimensional self-dual Galois representations have density zero by Rohrlich.

Unifying these results or obtaining asymptotical results on the automorphic side with fixed eigenvalue and increasing level, seems to be an interesting and hard problem.

2. Background

In this chapter we collect various background information on functoriality, results related to Weyl's Law, and some consequences of the Langlands-Shahidi method for irreducibility of induced representations that we will need (in Sections 2.2, 2.3, and 2.4, respectively). Note that further background on Laplacian eigenvalues and depth is in their respective chapters.

2.1 Basic Definitions and Notation

To fix the notation, let us review the definitions of the following reductive algebraic groups defined over $\mathbb{Q}$ with which we shall be mostly dealing. Let I_n be the $n \times n$ identity matrix and J_n the $n \times n$ matrix with 1's along the second diagonal.

$$GL_N = \{g | g \text{ is invertible}\}$$

$$SL_N = \{g | \det g = 1\}$$

$$SO_{2n+1} = \{g \in SL_{2n+1} | gM^t g = M\}, \text{ where } M = \begin{pmatrix} 0 & 0 & I_n \\ 0 & 1 & 0 \\ I_n & 0 & 0 \end{pmatrix}$$

$$Sp_{2n} = \{g \in SL_{2n} | gM^t g = M\}, \text{ where } M = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

$$SO_{2n} = \{g \in SL_{2n} | gM^t g = M\}, \text{ where } M = \begin{pmatrix} 0 & I_n \\ I_n & 0 \end{pmatrix}$$

$SO^*_{2n}(F) = SO^{*,\tau}_{2n}(F) = \{g \in SL_{2n}(F) | gM^t g = M\}$ for a field $F \supset \mathbb{Q}$ and $\tau \in F$ such that $-\tau$ is not a square in F, where $M = M_\tau = \begin{pmatrix} 0 & 0 & J_{n-1} \\ 0 & \Lambda_\tau & 0 \\ J_{n-1} & 0 & 0 \end{pmatrix}$ and $\Lambda_\tau = \begin{pmatrix} 1 & 0 \\ 0 & \tau \end{pmatrix}$

The group $SO^{*,\tau}_{2n}(F)$ is attached to a quadratic extension $F(\sqrt{-\tau})/F$, to which class field theory associates a quadratic character $\eta_{F(\sqrt{-\tau})/F}$.

All of the groups above are split, except for $SO^{*,\tau}_{2n}$, which is quasisplit but not split (when $-\tau$ is not a square).

Throughout, by a *quasisplit classical group*, we shall mean one of the groups SO_N, SO^*_{2n}, Sp_{2n} defined above.

For a number field F, we denote by $\mathbb{A}_F$ the ring of adeles of F. When $F = \mathbb{Q}$, we often just write $\mathbb{A}$ for $\mathbb{A}_\mathbb{Q}$. Denote the ring of integers of $\mathbb{Q}_p$ by $\mathcal{O}_p$.

We shall generally denote algebraic groups by boldface letters $\mathbf{G}, \mathbf{H}, \mathbf{M}$ and their F-points by G, H, M, etc.

Throughout the thesis, $\mathrm{Ind}_H^G \sigma$ denotes the non-normalized induction (unless explicitly stated otherwise). In Section 2.4 we shall define the normalized (twisted) parabolic induction $I(\nu, \sigma)$.

2.2 Langlands Functoriality and Reciprocity

Let us now review some theorems about Langlands correspondence and functoriality for GL_N and classical groups. We shall mostly follow [3] and [9].

Let F be a local or global field of characteristic zero. Let W_F be the Weil group, and when F is non-archimedean, let $W'_F = W_F \times SU(2)$ be the Weil-Deligne group. To provide a unified treatment of Langlands correspondence in the archimedean and

p-adic cases, denote by L_F the local Langlands group defined as W_F when F is archimedean and W'_F when F is non-archimedean.

Let $\mathbf{G}$ be a connected reductive group over F. Let $G^\vee$ be the (complex) Langlands dual group and ${}^L G = G^\vee \rtimes W_F$ the L-group of G.

Let us first take F to be local. Denote by $\Phi(G)$ the set of $G^\vee$-orbits of semisimple continuous L-homomorphisms $\varphi : L_F \to {}^L G$. Let $\Pi(G)$ be the set of equivalence classes of irreducible admissible representations π of $G = \mathbf{G}(F)$.

The local Langlands correspondence then conjectures that there is a reciprocity map from $\Pi(G)$ to $\Phi(G)$ with various natural properties, e.g., that it preserves L-functions. The fibers of the reciprocity map are called L-packets and are expected to be finite.

Suppose now that we have two reductive groups $\mathbf{G}_1$ and $\mathbf{G}_2$ and a homomorphism $f : {}^L G_1 \to {}^L G_2$. Each (irreducible admissible) representation π_1 of G_1 is expected to correspond to some $\varphi : L_F \to {}^L G_1$. Composing φ with f we obtain $f \circ \varphi : L_F \to {}^L G_2$, to which should correspond a representation π_2 of G_2. This gives us a conjectural map from $\Pi(G_1)$ to $\Pi(G_2)$, referred to as local Langlands functoriality.

Local Langlands correspondence and functoriality have been proved by Langlands [21] when $F = \mathbb{R}$ for all groups. When F is non-archimedean and $\mathbf{G} = GL_N$, reciprocity is due to Harris-Taylor [15] and Henniart [16]. Assuming the stabilization of the twisted trace formula for GL_N, Arthur [3] has recently proved reciprocity for classical groups $\mathbf{G}$ (only up to outer conjugacy by O_{2n} in the case of even orthogonal groups) and functoriality corresponding to natural (endoscopic) embeddings ${}^L G \hookrightarrow {}^L GL_N$. We shall discuss his results in more detail at the end of this section.

Take now a global field F. It is not known what extension L_F of the Weil group we need to take in order to expect Langlands reciprocity between representations $\varphi : L_F \to {}^L G$ and automorphic representations of $\mathbf{G}(\mathbb{A}_F)$ to hold. But we still

can consider conjectural global functoriality between automorphic representations of $\mathbf{G}_1(\mathbb{A}_F)$ and $\mathbf{G}_2(\mathbb{A}_F)$ as above.

Functoriality from classical groups to GL_N has been established by Cogdell, Kim, Piatetski-Shapiro, and Shahidi [8], [9] for generic automorphic representations. A precise description of the image of functoriality is given by the descent method of Ginzburg, Rallis, and Soudry [13]:

Theorem 2.2.1 *([9], [13]) Let* $\mathbf{G}$ *be a quasisplit classical group with an embedding* ${}^LG \hookrightarrow {}^LGL_N$ *and let* χ_G *and* R *be as in Table 2.1.*

Let π *be a globally generic cuspidal automorphic representation of* $\mathbf{G}(\mathbb{A}_F)$*. Then the functorial lift of* π *to an automorphic representation* Π *of* $GL_N(\mathbb{A}_F)$ *is self-dual with central character* $\omega_\Pi = \chi_G$*.* Π *is the normalized parabolic induction* $\Pi = \mathrm{Ind}(\Pi_1 \otimes \cdots \otimes \Pi_k) = \Pi_1 \boxplus \cdots \boxplus \Pi_k$*, where each* Π_i *is a self-dual cuspidal automorphic representation of* $GL_{N_i}(\mathbb{A}_F)$ *such that the partial L-function* $L^T(s, \Pi_i, R)$ *(with* T *a sufficiently large set of places of* F *containing all archimedean ones) has a pole at* $s = 1$ *and* $\Pi_i \not\cong \Pi_j$ *for* $i \neq j$*. Moreover, every such* Π *is a functorial lift of some* π*.*

Table 2.1.

Functoriality

$\mathbf{G}$	$\mathbf{G}^\vee$	N	R	χ_G
Sp_{2n}	SO_{2n+1}	$2n+1$	Sym^2	$\mathbf{1}$
SO_{2n+1}	Sp_{2n}	$2n$	$\bigwedge^2$	$\mathbf{1}$
SO_{2n}	SO_{2n}	$2n$	Sym^2	$\mathbf{1}$
SO^*_{2n}	SO_{2n}	$2n$	Sym^2	$\eta_{F(\sqrt{-\tau})/F}$

For all automorphic representations of classical groups, not only the generic ones, global functoriality was proved by Arthur [3] (again assuming the stabilization of

the twisted trace formula for GL_N and only up to outer conjugacy by O_{2n} for even orthogonal groups).

Take $\mathbf{G}$ to be a quasisplit classical group and let us now briefly review some of Arthur's results which we shall use, starting again with the local case:

Let $\widetilde{\Psi}^+(G)$ the set of (not necessarily irreducible) L-homomorphisms $\psi : L_F \times SU(2) \to {}^LG$, taken up to conjugacy by $\mathrm{Out}(G^\vee)$. Arthur then defines subsets $\widetilde{\Psi}_{\mathrm{sim}}(G) \subset \widetilde{\Psi}_2(G) \subset \widetilde{\Psi}(G) \subset \widetilde{\Psi}^+_{\mathrm{unit}}(G) \subset \widetilde{\Psi}^+(G)$ and $\widetilde{\Phi}_{\mathrm{bdd}}(G)$. Let $\widetilde{\Pi}_{\mathrm{unit}}(G)$ be the set of of unitary irreducible admissible representations of G (taken up to conjugacy by outer automorphisms of G) and $\widetilde{\Pi}_{\mathrm{temp}}(G)$ the set of those which are tempered. Finally, for $\psi \in \widetilde{\Psi}(G)$ we have the group $\mathcal{S}_\psi$. See Chapter 1 of [3] for precise definitions and details.

We then have the local classification of Theorem 1.5.1 in [3].

Theorem 2.2.2 *([3], 1.5.1) a) For every $\psi \in \widetilde{\Psi}(G)$, there is a finite subset $\widetilde{\Pi}_\psi$ of $\widetilde{\Pi}_{\mathrm{unit}}(G)$ equipped with a canonical mapping*

$$\pi \mapsto \langle \cdot, \pi \rangle, \qquad \pi \in \widetilde{\Pi}_\psi,$$

from $\widetilde{\Pi}_\psi$ to the group $\hat{\mathcal{S}}_\psi$ of characters on $\mathcal{S}_\psi$ such that $\langle \cdot, \pi \rangle = 1$ if G and π are unramified (relative to a suitable fixed compact open subgroup K).

b) If $\varphi = \psi$ belongs to the subset $\widetilde{\Phi}_{\mathrm{bdd}}(G)$ of parameters in $\widetilde{\Psi}(G)$ which are trivial on the factor $SU(2)$, the elements in $\widetilde{\Pi}_\varphi$ are tempered and the corresponding mapping from $\widetilde{\Pi}_\phi$ to $\hat{\mathcal{S}}_\psi$ is injective. Moreover, every element in $\widetilde{\Pi}_{\mathrm{temp}}(G)$ belongs to exactly one packet $\widetilde{\Pi}_\phi$. Finally, if F is non-archimedean, the mapping from $\widetilde{\Pi}_\varphi$ to $\hat{\mathcal{S}}_\psi$ is bijective.

To a parameter $\psi \in \widetilde{\Psi}^+(G)$ one can associate a parameter $\varphi_\psi \in \widetilde{\Phi}(G)$, given by

$$\varphi_\psi(w) = \psi(w, \begin{pmatrix} |w|^{1/2} & 0 \\ 0 & |w|^{-1/2} \end{pmatrix}.$$

Then $\widetilde{\Pi}_{\varphi_\psi} \subset \widetilde{\Pi}_\psi$ and for K-spherical representations we have the following proposition:

Proposition 2.2.3 *Let* $\mathbf{G}$ *be a split orthogonal or symplectic group and* K *a fixed maximal compact subgroup of* $\mathbf{G}(F)$*; if* F *is non-archimedean we take* $K = \mathbf{G}(\mathcal{O}_F)$*.*

Let $\psi \in \widetilde{\Psi}^{+}_{\mathrm{unit}}(G)$ *and assume that* $\pi \in \widetilde{\Pi}_{\psi}$ *has a non-zero* K*-fixed vector. Then* $\pi \in \widetilde{\Pi}_{\varphi_\psi}$*. Moreover, there is at most one such a representation* π *in* $\widetilde{\Pi}_{\psi}$*.*

Let us note that similar results were discussed by Shahidi [33].

Let us now discuss the global results, and so assume that F is global. We denote by $\mathcal{A}(G)$, $\mathcal{A}_2(G)$ and $A_{\mathrm{cusp}}(G)$ the sets of (unitary) automorphic representations of $\mathbf{G}(\mathbb{A}_F)$, respectively those that are in the discrete spectrum or cuspidal.

Let $\Psi_{\mathrm{sim}}(N) = \Psi_{\mathrm{sim}}(GL(N))$ be the set of formal tensor products $\psi = \mu \boxtimes \nu$, where $N = mn$, $\mu \in \mathcal{A}_{\mathrm{cusp}}(GL(m))$ and ν is the unique irreducible representation of $SU(2)$ of degree n. The parameter ψ corresponds to an isobaric sum

$$\Pi = \mu\left(\frac{n-1}{2}\right) \boxplus \mu\left(\frac{n-3}{2}\right) \boxplus \cdots \boxplus \mu\left(-\frac{n-1}{2}\right),$$

where $\mu(i) : x \mapsto \mu(x)|\det x|^i$. By a Theorem of Moeglin-Waldspurger, the set $\Psi_{\mathrm{sim}}(N)$ parametrizes the discrete spectrum $\mathcal{A}_2(GL(N))$.

Let $\widetilde{\Psi}_{\mathrm{sim}}(N)$ be the subset of self-dual elements in $\Psi_{\mathrm{sim}}(N)$ and let $\widetilde{\Psi}_2(N)$ be the set of formal unordered sums $\psi = \psi_1 \boxplus \cdots \boxplus \psi_r$ for distinct elements $\psi_i \in \widetilde{\Psi}_{\mathrm{sim}}(N_r)$, where $N = N_1 + \cdots + N_r$. Elements of $\widetilde{\Psi}_2(N)$ thus correspond to isobaric sums $\Pi_1 \boxplus \cdots \boxplus \Pi_r$ for $\Pi_i \in \mathcal{A}_2(GL(N_i))$.

Finally, let $\widetilde{\Psi}_2(G)$ be the set of those parameters $\psi \in \widetilde{\Psi}_2(N)$, which are expected to be lifts of automorphic representations from $\mathbf{G}(\mathbb{A}_F)$ (see [3], Section 1.4 for a precise definition).

We have a localization mapping $\psi \mapsto \psi_v$ from $\widetilde{\Psi}_2(G)$ to $\widetilde{\Psi}^{+}_{\mathrm{unit}}(G_v)$ and so we can define the global packet

$$\widetilde{\Pi}_{\psi} = \left\{ \bigotimes_v \pi_v | \pi_v \in \widetilde{\Pi}_{\psi_v}, \langle \cdot, \pi_v \rangle = 1 \text{ for almost all } v \right\}.$$

Let $\widetilde{\mathcal{H}}(G)$ be the product of the local symmetric Hecke algebras.

Theorem 2.2.4 *([3], 1.5.2) There is an $\widetilde{\mathcal{H}}(G)$-module isomorphism*

$$L^2_{\mathrm{disc}} \simeq \bigoplus_{\psi \in \widetilde{\Psi}_2(G)} \bigoplus_{\pi \in \widetilde{\Pi}_\psi(\varepsilon_\psi)} m_\psi \pi,$$

where m_ψ equals 1 or 2, $\varepsilon_\psi : \mathcal{S}_\psi \to \{\pm 1\}$ is an explicitly defined character and $\widetilde{\Pi}_\psi(\varepsilon_\psi)$ is the subset of elements of $\widetilde{\Pi}_\psi$ such that $\langle \cdot, \pi \rangle = \varepsilon_\psi$.

When G is not an even orthogonal group, m_ψ is always 1.

Finally, let us consider K-spherical representations in an A-packet $\widetilde{\Pi}_\psi$ for $K = K_\infty \times \prod K_p$ with $K_p = \mathbf{G}(\mathcal{O}_p)$ and a fixed maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ (to simplify notation a little, we take $\mathbf{G}$ split and $F = \mathbb{Q}$ here).

Lemma 2.2.5 *Let $\mathbf{G}$ be a split classical group, $F = \mathbb{Q}$ and $\psi \in \widetilde{\Psi}_2(G)$. Then there is at most one representation $\pi \in \widetilde{\Pi}_\psi$ such that $\pi^K \neq 0$ (where K is the maximal compact subgroup as above).*

Proof Let $\pi = \otimes \pi_v$. Each π_v is K_v-spherical, and so by Proposition 2.2.3, there is at most one such π_v. Hence there is at most one global representation π. □

Note that we need general Langlands correspondence and functoriality from [3] only in the last chapter where we prove a lower bound on the number of self-dual representations. In the rest of the thesis we could just work with the generic functoriality (which is available unconditionally). The analogue of Proposition 4.2.3 is in fact easier to prove only for generic representations than in general.

2.3 Weyl's Law

Let $\mathbf{G}$ be a reductive group over $\mathbb{Q}$. For a prime p, take a compact open subgroup K_p of $\mathbf{G}(\mathbb{Q}_p)$, also let K_∞ be a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$. This gives us a compact subgroup $K = K_\infty \times \prod K_p$ of $\mathbf{G}(\mathbb{A})$. Finally, denote by $A_{\mathbf{G}}$ the maximal

split torus in the center of $\mathbf{G}$ and let $A_{\mathbb{R}} = A_{\mathbf{G}}(\mathbb{R})^0$ be the connected component of the $\mathbb{R}$-points of $A_{\mathbf{G}}$.

Weyl's Law conjecturally gives asymptotic results for the number of cuspidal automorphic representations of $\mathbf{G}(\mathbb{A})$. For $\lambda > 0$, define $N_G^K(\lambda) = \sum_{\lambda(\pi)\leq\lambda} \dim \pi^K$, where the sum is over all cuspidal automorphic representations $\pi = \otimes\pi_v$ of $\mathbf{G}(\mathbb{A})$ such that the Laplacian eigenvalue $\lambda(\pi)$ of the restriction of π_∞ to $\mathbf{G}^1(\mathbb{R})$ is at most λ (see Section 3.2 for the definition of the Laplacian eigenvalue). Here $\mathbf{G}^1 = \{g \in \mathbf{G} |\ |\chi(g)| = 1$ for all $\chi \in X^*(\mathbf{G})\}$.

Weyl's Law then asserts that $N_G^K(\lambda) = c\lambda^{d/2} + o(\lambda^{d/2})$ as $\lambda \to \infty$ for an explicit constant c, where d is the dimension of the locally symmetric space

$$X = X_G^K = A_{\mathbb{R}}\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})/K.$$

In this form, Weyl's Law was proved for $\mathbf{G} = GL_N$ by Müller [27] and then by Lapid-Müller [22] with an improved error term:

Theorem 2.3.1 *[27] Let $\mathbf{G} = GL_N$, let $K_p = K_p(m)$ be the principal congruence subgroup of level m (i.e., the subgroup consisting of matrices congruent to the identity modulo p^m) and $K_\infty = O(N)$. Then*

$$N_{GL(N)}^K(\lambda) = \frac{\mathrm{vol}(X)}{(4\pi)^{d/2}\mathbf{\Gamma}(d/2+1)}\lambda^{d/2} + o(\lambda^{d/2}),$$

where $\mathbf{\Gamma}$ is the standard gamma function.

When $\mathbf{G}$ is a split adjoint semisimple group, Weyl's Law was proved by Lindenstrauss-Venkatesh [23]. It seems that a similar proof should work for a general split semisimple group, but this has not appeared in literature yet. At the very least, one has weak Weyl's Law:

Theorem 2.3.2 *There are constants c_1 and c_2 such that for all sufficiently large λ, one has $c_1\lambda^{d/2} < N_G^K(\lambda) < c_2\lambda^{d/2}$.*

When $\mathbf{G}$ is a split simply connected semisimple group, Theorem 2.3.2 was proved by Labesse-Müller [20].

On a general group the full Weyl's Law hasn't been proved yet, but we have the following upper bound due to Donnelly [10].

Let us first review the definition of an arithmetic subgroup: A subgroup H of a Lie group G is arithmetic if there exists $k \in \mathbb{N}$ and a faithful representation $\rho : G \to GL_k(\mathbb{C})$ such that $\rho(H)$ is commensurable with $\rho(G) \cap GL_k(\mathbb{Z})$. This condition does not depend on the choice of k and ρ.

Theorem 2.3.3 *[10] Let G be a (non-compact) semisimple Lie group, K a maximal compact subgroup of G and Γ an arithmetic subgroup of G. Let $N(\lambda)$ be the number of linearly independent Laplacian eigenfunctions in $L^2_{cusp}(K\backslash G/\Gamma)$ with eigenvalue less than λ. Then*

$$\limsup_{\lambda\to\infty} \frac{N(\lambda)}{\lambda^{d/2}} \leq \frac{\mathrm{vol}(K\backslash G/\Gamma)}{(4\pi)^{d/2}\mathbf{\Gamma}(d/2+1)},$$

where d is the dimension of $K\backslash G/\Gamma$.

The dimensions of the symmetric spaces are given in Table 2.2.

Table 2.2.

Dimensions

G	$\dim G$	$\dim K$	$d_G = \dim X_G$
$SL_{2n+1}(\mathbb{R})$	$4n^2+4n$	$2n^2+n$	$2n^2+3n$
$SL_{2n}(\mathbb{R})$	$4n^2-1$	$2n^2-n$	$2n^2+n-1$
$Sp_{2n}(\mathbb{R})$	$2n^2+n$	n^2	n^2+n
$SO_{2n+1}(\mathbb{R})$	$2n^2+n$	n^2	n^2+n
$SO_{2n}(\mathbb{R})$	$2n^2-n$	n^2-n	n^2
$SO^*_{2n}(\mathbb{R})$	$2n^2-n$	n^2	n^2-n

We shall need an adelic reformulation of Donnelly's result:

Proposition 2.3.4 *Let* $\mathbf{G}$ *be a connected reductive group such that* $\mathbf{G}^1(\mathbb{R})$ *is not compact. Let* K *be an arithmetic subgroup of* $\mathbf{G}(\mathbb{A})$ *and* $X = A_{\mathbb{R}}\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})/K$.

Then for λ *large enough we have*

$$N_G^K(\lambda) \leq \frac{\mathrm{vol}(X)}{(4\pi)^{d/2}\mathbf{\Gamma}(d/2+1)}\lambda^{d/2},$$

where d *is the dimension of* X.

Here we extend the definition of an arithmetic subgroup to the adelic setting as follows: Let $K_f = \prod_{p<\infty} K_p$. By the strong approximation theorem we have that $A_{\mathbb{R}}\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})/\mathbf{G}(\mathbb{R})K_f$ is finite. Denote by $x_1 = 1, x_2, \ldots, x_l$ the double coset representatives in $\mathbf{G}(\mathbb{A}_f)$ and let $\Gamma_i = (\mathbf{G}(\mathbb{Q}) \cdot x_i K_f x_i^{-1}) \cap \mathbf{G}^1(\mathbb{R})$. We define K to be arithmetic if each of these Γ_i is an arithmetic subgroup of $\mathbf{G}(\mathbb{R})$ (as defined before Theorem 2.3.3).

Proof We argue similarly as on page 134 of [22]. Let Γ_i be as above. Since K is arithmetic, all of these Γ_i are arithmetic subgroups of $\mathbf{G}(\mathbb{R})$ by definition.

We obtain an isomorphism of $\mathbf{G}(\mathbb{R})$-spaces

$$A_{\mathbb{R}}\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})/K_\infty K_f \simeq \bigsqcup_i \Gamma_i\backslash\mathbf{G}^1(\mathbb{R})/K_\infty.$$

Automorphic representations correspond to automorphic forms on $\Gamma_i\backslash\mathbf{G}^1(\mathbb{R})/K_\infty$; denote by $N_i(\lambda)$ the number of linearly independent Laplacian eigenfunctions in the space $L^2_{cusp}(\Gamma_i\backslash\mathbf{G}^1(\mathbb{R})/K_\infty)$ with eigenvalue less than λ. Then $N_G^K(\lambda) = \sum_i N_i(\lambda)$.

Using Donnelly's upper bound for each $N_i(\lambda)$, we obtain the desired estimate for $N_G^K(\lambda)$. □

Note that the assumption that $\mathbf{G}^1(\mathbb{R})$ is not compact is satisfied for all of the groups with which we shall be dealing, except for non-split SO_2^*.

2.4 L-Functions and Irreducibility of Induced Representations

In this Section we review some results concerning irreducibility of parabolically induced representations, which follow from the Langlands-Shahidi method. We shall use these results in Section 4.5. Our main reference is [32].

Let F be a non-archimedean field of characteristic 0, $\mathbf{G}$ a quasisplit connected reductive group over F and $G = \mathbf{G}(F)$.

The theory works more generally as well, but for our purposes, let us assume that $P = MN$ is a standard maximal parabolic subgroup of G. Let A be the split component of M, $\mathfrak{a}^* = X^*(M) \otimes \mathbb{R}$ and $\mathfrak{a} = \mathrm{Hom}(X^*(M), \mathbb{R})$. Define a map $H_M : M \to \mathfrak{a}$ by $\exp\langle \chi, H_M(m)\rangle = |\chi(m)|_F$. Denote by ρ_P half of the sum of roots in $\mathrm{Lie}(N)$ and by α the unique simple root of A in $\mathrm{Lie}(N)$. Finally, set $\tilde{\alpha} = \langle \rho_P, \alpha\rangle^{-1}\rho_P \in \mathfrak{a}^*$.

Let σ be an irreducible unitary generic (admissible) representation of M. For $\nu \in \mathfrak{a}_{\mathbb{C}}^*$ denote by $I(\nu, \sigma)$ the induced representation $\mathrm{Ind}_P^G \, \sigma \otimes \exp\langle \nu + \rho_P, H_M(\cdot)\rangle \otimes \mathbf{1}$.

To the parabolic P we can associate representations r_i with highest weight vectors given by the diagrams starting on page 183 of [32]. Using the Langlands-Shahidi method, one can then define the local coefficients $C_\chi(s\tilde{\alpha}, \sigma, w_0)$ and L-functions $L(s, \sigma, r_i)$.

We have the following theorem:

Theorem 2.4.1 *([31], 3.3.1 and [32], 8.4.9, 8.5.1) Assume that σ is an irreducible unitary generic supercuspidal representation of M. Then*

$$C_\chi(s\tilde{\alpha}, \sigma, w_0) \sim L(1-s, \tilde{\sigma}, r_1)L(1-2s, \tilde{\sigma}, r_2)/L(s, \sigma, r_1)L(2s, \sigma, r_2),$$

where $\sim$ means equality up to a monomial in q^{-s}. The following are equivalent:

a) $L(s, \sigma, r_i)$ has a pole at $s = 0$ for $i = 1$ or 2 and only for one of them.

b) $I(0, \sigma)$ is irreducible and σ is ramified, i.e., $w_0(\sigma) \simeq \sigma$.

Moreover, assume that these conditions hold and let $i = 1$ or 2 be such that $L(s, \sigma, r_i)$ has a pole at $s = 0$. Then for $s > 0$, we have that $I(s\tilde{\alpha}, \sigma)$ is reducible only when $s = 1/i$, for all other s it is irreducible.

If σ is ramified and $I(0, \sigma)$ is reducible, then all $I(s\tilde{\alpha}, \sigma)$ with $s > 0$ are irreducible.

If σ is unramified, then all $I(s\tilde{\alpha}, \sigma)$ with $s \geq 0$ are irreducible.

Let us now apply this theorem to prove a lemma which we shall need in Section 4.5.

Lemma 2.4.2 *Let* $\mathbf{G} = SO_{2n+1}$ *and* $P = MN$ *be standard maximal parabolic with Levi* $\mathbf{M} = GL_1 \times SO_{2n-1}$. *Let* σ *be an irreducible unitary generic supercuspidal representation of* $SO_{2n-1}(F)$ *and* χ *a unitary character of* $F^\times$ *such that* $\chi^2 \neq \mathbf{1}$. *Then the induced representation* $\mathrm{Ind}_P^G\, \sigma \otimes \chi$ *is irreducible.*

Proof Since $\chi^2 \neq \mathbf{1}$, we have that $\sigma \otimes \chi$ is unramified in the sense of Theorem 2.4.1. By this theorem we then see that all $I(s\tilde{\alpha}, \sigma \otimes \chi)$ are irreducible. In particular, $I(\tilde{\alpha}/2, \sigma \otimes \chi)$ is irreducible.

We have $\alpha = e_1 - e_2$ and $\rho_P = e_2/2$. Hence $\langle \rho_P, \alpha \rangle = -1/2$, and so $\tilde{\alpha} = -2\rho_p$. Hence we see that $I(\tilde{\alpha}/2, \sigma \otimes \chi) = I(-\rho_P, \sigma \otimes \chi)$ is the non-normalized induction $\mathrm{Ind}_P^G\, \sigma \otimes \chi$, the irreducibility of which we wanted to prove. □

3. Laplacian Eigenvalues

In this chapter we shall study the behavior of Laplacian eigenvalues under functoriality. This is given by Theorem 3.3.2, which we will later use in the proofs in Chapters 5 and 6.

Laplacian eigenvalues are obtained by evaluating the infinitesimal character on the Casimir element. Hence we start by discussing the Casimir elements in Section 3.1 and then consider the infinitesimal characters in Section 3.2. Finally, we prove Theorem 3.3.2 in Section 3.3.

Our main reference is [19], Chapter V.5.

Throughout this chapter, all groups and representations are real, unless stated otherwise.

3.1 Casimir Element

In this Section, we review the definition of the Killing form and Casimir element, following [19] (mostly Chapter V).

Let $\mathfrak{g}$ be a Lie algebra (viewed as a Lie subalgebra of $\mathrm{Mat}_{m\times m}(\mathbb{C})$ for some m). For $X, Y \in \mathfrak{g}$, we define the Killing form as $B(X,Y) = \mathrm{Tr}(\mathrm{ad}\ X\ \mathrm{ad}\ Y)$. It is a symmetric bilinear form on $\mathfrak{g}$. We also have the bilinear form $C(X,Y) = \mathrm{Tr}(XY)$. When $\mathfrak{g}$ is a simple complex Lie algebra, $B(X,Y) = b_{\mathfrak{g}}C(X,Y)$ for some constant $b_{\mathfrak{g}}$.

We shall be dealing mostly with the Lie algebras $\mathfrak{sl}_N(\mathbb{C})$, $\mathfrak{sp}_{2n}(\mathbb{C})$, $\mathfrak{so}_N(\mathbb{C})$, and $\mathfrak{so}^*_{2n}(\mathbb{C})$. For each of them, the Killing form is a multiple of $C(X,Y)$ and the corresponding constants are $b_{\mathfrak{sl}(N)} = 2N$, $b_{\mathfrak{sp}(2n)} = 2n+2$, and $b_{\mathfrak{so}(N)} = b_{\mathfrak{so}^*(N)} = N-2$ (see eg. [11], Ex. 14.36).

Now, let $\mathfrak{g}$ be a real Lie algebra and $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ its complexification. Let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$ (and $\mathfrak{h}_{\mathbb{C}}$ its complexification). Let Φ be the set of roots of $\mathfrak{g}_{\mathbb{C}}$ with respect to $\mathfrak{h}_{\mathbb{C}}$.

Choose an orthonormal basis $H_1, \ldots, H_n$ of $\mathfrak{h}_{\mathbb{C}}$ with respect to the Killing form B on $\mathfrak{g}_{\mathbb{C}}$. Also choose root vectors E_α so that $B(E_\alpha, E_{-\alpha}) = 1$ for all $\alpha \in \Phi$. Then the Casimir element of $\mathfrak{g}_{\mathbb{C}}$ is $\Omega = \sum_{i=1}^{n} H_i^2 + \sum_{\alpha \in \Phi} E_\alpha E_{-\alpha}$ as an element of the center $Z(\mathfrak{g}_{\mathbb{C}})$ of the universal enveloping algebra.

We have the Harish-Chandra isomorphism $\gamma : Z(\mathfrak{g}_{\mathbb{C}}) \to \mathcal{H}^W$, where $\mathcal{H}$ is the universal enveloping algebra of $\mathfrak{h}_{\mathbb{C}}$ and $\mathcal{H}^W$ are its Weyl group invariants. For more details, see [19], Ch. V.5.

To compute the Laplacian eigenvalues of our representations, we shall need the images of the Casimir element under the Harish-Chandra isomorphism. In general, it is $\gamma(\Omega) = \sum H_i^2 - |\delta|^2$, where δ is half of the sum of positive roots.

3.2 Infinitesimal Character

Let G be a Lie group and let $\mathfrak{g}$, $\mathfrak{h}$, $\mathfrak{g}_{\mathbb{C}}$, and $\mathfrak{h}_{\mathbb{C}}$ be the corresponding Lie algebras as above.

For $\mu \in \mathfrak{h}_{\mathbb{C}}^*$ define $\chi_\mu(Z) = \mu(\gamma(Z))$ for $Z \in Z(\mathfrak{g}_{\mathbb{C}})$. Note that $\gamma(Z) \in U(\mathfrak{h}_{\mathbb{C}}) = \mathcal{H}$, and so it makes sense to evaluate μ on $\gamma(Z)$ – here we are of course extending μ to a character of the universal enveloping algebra $\mathcal{H}$. It follows from the Harish-Chandra isomorphism that all characters of $Z(\mathfrak{g}_{\mathbb{C}})$ are of this form.

Let π be an irreducible admissible representation of G. From π we can obtain a representation of $Z(\mathfrak{g}_{\mathbb{C}})$. This representation is then a character, called the infinitesimal character of π and denoted by χ_π. This character is of the form $\chi_\pi = \chi_\mu$ for some $\mu \in \mathfrak{h}_{\mathbb{C}}^*$ as above. This μ is uniquely determined up to the action of W and it is sometimes also called the infinitesimal character of π.

The Laplacian eigenvalue of π is then obtained by evaluating the infinitesimal character on the Casimir element, i.e., $\chi_\pi(\Omega)$.

3.3 Behavior under Functoriality

To see what happens to the Laplacian eigenvalue under functoriality, we need to study what happens to the infinitesimal character.

Let $W_{\mathbb{R}}$ be the Weil group. By the local Langlands correspondence, to each π corresponds a parameter $\varphi : W_{\mathbb{R}} \to {}^L G$, where ${}^L G$ is the Langlands dual group. By conjugating φ if necessary, we may assume that $\varphi(W_{\mathbb{R}}) \subset N({}^L H^0)$, where H is a maximal torus in G. Then φ is of the form $\varphi(z) = z^{\mu}\overline{z}^{\mu'}$ for $z \in \mathbb{C}^{\times} \subset W_{\mathbb{R}}$, where $\mu, \mu' \in X_*(H^{\vee}) \otimes \mathbb{C} = X^*(H) \otimes \mathbb{C}$ are such that $\mu - \mu' \in X^*(H)$. Here $\varphi(z) = z^{\mu}\overline{z}^{\mu'}$ stands for $\varphi(e^s) = \exp(s\mu + \overline{s}\mu') \in {}^L H^0$ for $s \in \mathbb{C}$.

By definition we have $X_*(H^{\vee}) = X^*(H)$, and so we can view μ as an element of $X^*(H) \otimes \mathbb{C}$ and attach to it a character χ_μ of $Z(\mathfrak{g}_{\mathbb{C}})$ via the Harish-Chandra homomorphism. If φ is the Langlands parameter of π, then χ_μ is the infinitesimal character of π. (See for example [1] for details.)

In the rest of this section, let $G = \mathbf{G}(\mathbb{R})$ be a quasisplit classical group. Let Π be a self-dual irreducible admissible representation of $GL_N(\mathbb{R})$ which is a functorial lift of an irreducible admissible representation π of G. The Laplacian eigenvalue of Π is determined from the infinitesimal character of the restriction of Π to $SL_N(\mathbb{R})$, and so we will be dealing with SL and $\mathfrak{sl}$.

Let $\varphi : W_{\mathbb{R}} \to {}^L G$ be the Langlands parameter of π. The Langlands parameter of Π is then given by composition with the inclusion ${}^L G \hookrightarrow {}^L SL_N$.

Proposition 3.3.1 *Let H be a maximal torus of G and T a maximal torus of $SL_N(\mathbb{R})$ such that $H^{\vee} \hookrightarrow T^{\vee}$. This induces an inclusion $\iota : X^*(H) \otimes \mathbb{C} = X_*(H^{\vee}) \otimes \mathbb{C} \hookrightarrow X_*(T^{\vee}) \otimes \mathbb{C} = X^*(T) \otimes \mathbb{C}$.*

Then the infinitesimal character of Π is $\iota(\mu)$, where μ is the infinitesimal character of π.

Proof Immediately follows from the discussion at the beginning of this section. $\square$

Theorem 3.3.2 *Let λ and Λ be the Laplacian eigenvalues of π and Π, respectively. Then there are constants $c_G > 0$ and d_G such that $\Lambda = c_G\lambda + d_G$.*

We have $c_{Sp(2n)} = \frac{2n+2}{2n+1}$, $c_{SO(2n)} = c_{SO^*(2n)} = \frac{n-1}{n}$, $c_{SO(2n+1)} = \frac{2n-1}{2n}$.

Proof To prove the theorem, we evaluate the infinitesimal characters on Casimir elements using Proposition 3.3.1.

Denote by E_{ij} the matrix with 1 at the position (i, j) and 0 elsewhere and denote E_{ii} also by E_i. Let $E_i^* \in \mathfrak{h}^*$ be the corresponding dual element. For $A = \sum a_i E_i \in \mathfrak{h}_{\mathbb{C}}$, we define $A^* = \sum a_i E_i^* \in \mathfrak{h}_{\mathbb{C}}^*$.

For $\mathfrak{g} = \mathfrak{sl}_N(\mathbb{R})$, let $\mathfrak{a}$ be the subalgebra of diagonal matrices with trace zero. Extend the set $E_i - E_{n+i}$ to a basis of $\mathfrak{a}_{\mathbb{C}}$ (where $N = 2n + \varepsilon$ with $\varepsilon = 0, 1$). We have $B(E_i - E_{n+i}, E_j - E_{n+j}) = 2N\mathrm{Tr}((E_i - E_{n+i})(E_j - E_{n+j})) = 2N\delta_{ij}$ (where δ_{ij} is Kronecker delta). Hence we can extend $A_i = \frac{1}{\sqrt{2N}}(E_i - E_{n+i})$ to an orthonormal basis of $\mathfrak{a}_{\mathbb{C}}$ with respect to the Killing form B.

Take $G = Sp_{2n}(\mathbb{R})$. Take $\mathfrak{h}_{\mathbb{C}}$ to be the subalgebra of diagonal matrices in $\mathfrak{g}$; we can choose $b_i = E_i - E_{n+i}$ as a basis of $\mathfrak{h}_{\mathbb{C}}$. We have $B(b_i, b_j) = (2n + 2)\mathrm{Tr}(b_i b_j) = 2(2n + 2)\delta_{ij}$. Thus $H_i = \frac{1}{\sqrt{2(2n+2)}} b_i$ form an orthonormal basis of $\mathfrak{h}_{\mathbb{C}}$ with respect to the Killing form B.

Let π be a representation of G with infinitesimal character $\mu = (\mu_1, \ldots, \mu_n) \in \mathfrak{h}_{\mathbb{C}}^*$ with respect to the basis $b_i^* = E_i^* - E_{n+i}^*$, i.e., $\mu = \sum \mu_i b_i^*$. We have $\gamma(\Omega) = \sum H_i^2 - |\delta_{Sp}|^2$ and $b_i^*(H_j) = \frac{2}{\sqrt{2(2n+2)}}\delta_{ij}$, and so we see that

$$\mu(\gamma(\Omega)) = \sum \mu_i^2 b_i^*(H_i)^2 - |\delta_{Sp}|^2 = \frac{2}{2n+2}\sum \mu_i^2 - |\delta_{Sp}|^2.$$

Let Π be the restriction of the functorial lift of π to $SL_{2n+1}(\mathbb{R})$. The infinitesimal character of Π is then $\iota(\mu) = (\mu_1, \ldots, \mu_n, -\mu_1, \ldots, -\mu_n) \in \mathfrak{a}_{\mathbb{C}}^*$. Note that $\iota(\mu) = \sum \mu_i (E_i - E_{n+i})^*$. Thus the Laplacian eigenvalue

$$\iota(\mu)(\gamma(\Omega)) = \sum \mu_i^2 (E_i - E_{n+i})^*(A_i)^2 - |\delta_{SL}|^2 = \frac{2}{2n+1}\sum \mu_i^2 - |\delta_{SL}|^2.$$

We see that $c_{Sp(2n)} = \frac{2n+2}{2n+1}$.

Take $G = SO_{2n}(\mathbb{R})$. This is entirely analogous to the case of Sp_{2n}, the only differences are that $B(b_i, b_j) = (2n-2)\mathrm{Tr}(b_i b_j)$ and that π lifts to $SL_{2n}(\mathbb{R})$, which leads to $c_{SO(2n)} = \frac{2n-2}{2n} = \frac{n-1}{n}$.

The case of $G = SO_{2n+1}(\mathbb{R})$ is also analogous, we get $c_{SO(2n+1)} = \frac{2n-1}{2n}$.

Let us finally discuss the case when $G = SO^*_{2n}(\mathbb{R})$ in more detail. Take G and $\mathfrak{g}$ to be defined with respect to the matrix $M = (m_{ij})$ with $m_{ij} = 1$ for $i + j = 2n$, $1 \leq i \leq n-1$, or $n \leq i = j \leq n+1$, and $m_{ij} = 0$ otherwise. Choose the Cartan algebra $\mathfrak{h}$ to be the span of $b_j = E_j - E_{2n-j}$ for $1 \leq j \leq n-1$, and $b_n = E_{n,n+1} - E_{n+1,n}$. Note that in this case the Killing form is not positive definite on $\mathfrak{h}$, as we have $B(b_n, b_n) = -2$. Setting $H_j = \frac{1}{\sqrt{2(2n-2)}} b_j$ we get that

$$\gamma(\Omega) = \sum_{j=1}^{n-1} H_j^2 - H_n^2 - |\delta_{SO^*}|^2.$$

Let π be a representation of G with infinitesimal character $\mu = (\mu_1, \ldots, \mu_n) \in \mathfrak{h}^*_{\mathbb{C}}$ with respect to the basis b_j^*. Then

$$\mu(\gamma(\Omega)) = \sum_{j \leq n-1} \mu_j^2 b_j^*(H_j)^2 - \mu_n^2 b_n^*(H_n)^2 - |\delta_{SO^*}|^2 = \frac{2}{2n-2}\Big(\sum_{j \leq n-1} \mu_j^2 - \mu_n^2\Big) - |\delta_{SO^*}|^2.$$

Let Π be the restriction of the functorial lift of π to $SL_{2n}(\mathbb{R})$ and let $\varphi : W_{\mathbb{R}} \to {}^L G$ be the Langlands parameter of π. Viewing μ as a cocharacter of $H^\vee$ we get that $\mu(\mathbb{C}^\times)$ is not in the torus ${}^L T^0$ – we have to conjugate the part corresponding to H_n, originally of the form $\begin{pmatrix} \cos\mu_n & \sin\mu_n \\ -\sin\mu_n & \cos\mu_n \end{pmatrix}$, which we conjugate to $\begin{pmatrix} e^{i\mu_n} & 0 \\ 0 & e^{-i\mu_n} \end{pmatrix}$ (here $i = \sqrt{-1}$). This gives us that the infinitesimal character of Π is then $\iota(\mu) = (\mu_1, \ldots, i\mu_n, -i\mu_n, \ldots, -\mu_1) \in \mathfrak{a}^*_{\mathbb{C}}$. Hence

$$\iota(\mu)(\gamma(\Omega)) = \frac{2}{2n}\Big(\sum \mu_j^2 - \mu_n^2\Big) - |\delta_{SL}|^2.$$

We conclude that $c_{SO^*(2n)} = \frac{2n-2}{2n} = \frac{n-1}{n}$.

This concludes the proof of Theorem 3.3.2. Note that one can also easily compute the constants d_G by calculating the various values $|\delta|^2$. $\square$

4. Depth

In this chapter we shall study representations of p-adic groups, in particular properties of their depth. A powerful tool for that is provided by Bruhat-Tits theory. We first briefly review some aspects of it in Section 4.1 and then, in Section 4.2, we prove the results we need for counting self-dual representation. In Section 4.3 we use the local descent to show a partial result towards depth preservation for split classical groups. However, depth need not be preserved in the case of wildly ramified groups, as we note in Section 4.4. The last Section, 4.5, is devoted to the study of the restriction of certain representations of SO_{2n} to SO_{2n-1}, which is one of the steps needed in trying to extend the results of Section 4.3 to a proof of full depth preservation, not just one inequality.

Throughout this chapter, all groups and representations are p-adic, unless stated otherwise.

4.1 Background

Throughout this chapter, let F be a local non-archimedean field (of characteristic zero) and G a connected reductive group over F. We shall study irreducible admissible representations of $G(F)$, their depth and behaviour under functoriality, so let's first review some background on Bruhat-Tits theory, Moy-Prasad filtrations, and depth. The basic references that we shall follow are [34], [35], and [26].

Let S be a maximal F-split torus of G, $N(S)$ the normalizer of S in G, and $Z(S)$ the centralizer of S in G. Let C be the maximal F-split torus contained in the center of G, we have $C \subset S$. Let $(\Phi, R, \Phi^\vee, R^\vee)$ be the relative root system, let W be its Weyl group and $\widetilde{W}$ the affine Weyl group, an extension of W by a free abelian group of rank $\dim S$. The (restricted) apartment $A = A(G, S, F)$ is defined as an affine

space under $(X_*(S)\otimes\mathbb{R})/(X_*(C)\otimes\mathbb{R})$, it is equipped by an action of $\widetilde{W}$ and by a corresponding action of $N(S)$. We denote both of these action by ν. See [34], 1.2 for details. Note that [34] works with the extended apartment, but the reduced one is somewhat more convenient for our purposes.

We also have the set of affine roots Φ_{af}, being the set of certain affine functions $\alpha = a+\gamma$ for $a\in\Phi$ and $\gamma\in\mathbb{R}$. See [34], 1.6.

For $a\in\Phi$ we have the root subgroup U_a; for an affine function $\alpha = a+r$, $r\in\mathbb{R}$ one defines a subgroup X_α of U_a as the subset of elements u satisfying $u=1$ or $\alpha(a,u)\geq\alpha$, where $\alpha(a,u)$ is a certain affine function, defined in [34], 1.4.

For an affine function $\alpha = a+r$ with $a\in\Phi$, we denote $A_\alpha = \alpha^{-1}([0,\infty))$, its boundary is $\partial A_\alpha = \alpha^{-1}(0)$, and r_α is the affine reflection whose vector part is the reflection r_a and whose fixed hyperplane is ∂A_α. When $\alpha\in\Phi_{\mathrm{af}}$, we call A_α the half-apartments and ∂A_α the walls. The chambers are the connected components of the complement in A of the walls (cf. [34], 1.7). A chamber is a polysimplex (i.e., a product of simplices); by a facet of A we mean just a facet of some chamber (viewed as a polysimplex).

The (restricted) building $\mathcal{B} = \mathcal{B}(G,F)$ is defined as the unique set equipped with a left G-action such that $\mathcal{B} = \bigcup_{g\in G} gA$, the group $N(S)$ stabilizes A and acts on it by ν, and for every affine root α, the group X_α fixes the half-apartment A_α (cf. [34], 2.1).

Fix a chamber $\mathcal{C}$. Then each element of A is $N(S)$-conjugate to an element of the closure $\bar{\mathcal{C}}$. Hence every element of $\mathcal{B}$ is G-conjugate to an element of the closure $\bar{\mathcal{C}}$.

For a facet $\mathcal{F}$, one can define a corresponding parahoric subgroup $G_{\mathcal{F}}$. The definition is somewhat involved ([35], 2.2.4), but roughly speaking, the parahoric subgroup is essentially the stabilizer of $\mathcal{F}$ in G. One can also define the parahoric subgroup G_x for $x\in\mathcal{B}$; if x is a "generic" point of a facet $\mathcal{F}$, then $G_{\mathcal{F}} = G_x$.

We have the Moy-Prasad subgroups which are used to define the depth of admissible representations [26]. First, denote by Z_n the group of all $z\in Z(S)$ such that

$\omega(\chi(z)-1) \geq n$ for all characters χ of $Z(S)$ (ω is the additive valuation on F). For $x \in \mathcal{B}$ and $r \geq 0$, we define $G_{x,r}$ to be the intersection of G_x with the subgroup generated by Z_n for $n \geq r$ and by X_α for all α satisfying $\alpha(x) \geq r$. Denote by $G_{x,r+}$ the union of all $G_{x,s}$ with $s > r$.

Let π be an irreducible admissible representation of G. The depth of π is defined as depth $\pi = \inf\{r \geq 0 | \exists x \in \mathcal{B}$ such that $\pi^{G_{x,r+}} \neq 0\}$.

One also defines the depth of a Weil-Deligne representation $\varphi : W'_F \to {}^L G$ as $\inf\{r \mid \varphi(I_s) = 1$ for all $s > r\}$, where I_s, $s \in \mathbb{Q}$, is the filtration of the inertia subgroup I of $W_F \subset W'_F$.

We have the following fundamental properties of depth:

- depth π is a rational number and the infimum is achieved for some $x \in \mathcal{B}$.
- The depth is preserved by parabolic induction and the Jacquet functor. Namely, let M be a Levi of G and σ an irreducible admissible representation of M. If π is an irreducible subquotient of the normalized induction $\mathrm{Ind}_M^G \sigma$, then depth σ = depth π (and similarly for the Jacquet functor).
- Depth preservation: Assume that $\mathbf{G}$ is unramified or tamely ramified. If $\varphi : W'_F \to {}^L G$ is the Langlands parameter of a representation π, then it is conjectured that depth π = depth ϕ. J.-K. Yu [35] has proved this for GL_N and for tamely ramified tori.

As we note in Section 4.4, depth need not be preserved in the case of wildly ramified groups. This has also been observed by Gross-Reeder [14]; Reeder-Yu [28] recently provided more examples of this phenomenon.

4.2 Bounds on Depth

We will use the following result on uniform admissibility, due originally to Howe when $\mathbf{G} = GL_N$, and then to Bernstein for general $\mathbf{G}$ (with a simpler proof) – for part b) see [5], Theorem 1; part a) then follows as in [6], Corollary 4.15.

Theorem 4.2.1 *([5], [6]) Let* $\mathbf{G}$ *be a reductive group over a non-archimedean local field* F*. Let* K *be an open compact subgroup of* $G = \mathbf{G}(F)$*. Then:*

a) There are finitely many supercuspidal representations σ *of* G *such that* $\sigma^K \neq 0$*.*

b) There is an $n \in \mathbb{N}$ *(depending only on* G *and* K*) such that, for all irreducible admissible representations* π *of* G*, we have* $\dim \pi^K \leq n$*.*

Both parts of Theorem 4.2.1 will be useful for us in the thesis. We shall shortly use the first part to prove part b) of Proposition 4.2.2. The second part of the theorem will play an important role in the arguments of Chapter 5, where it will enable us to uniformly bound the dimensions of the spaces of K-fixed vectors of the self-dual representations which we are counting.

Proposition 4.2.2 *Let* $\mathbf{G}$ *be a reductive group over a non-archimedean local field* F*.*

a) For each $r \geq 0$ *there is an open compact subgroup* K *of* $G = \mathbf{G}(F)$ *such that if* $\text{depth } \pi \leq r$*, then* $\pi^K \neq 0$*. When* $\mathbf{G}$ *is split and almost simple, we can take* K *to be the principal congruence subgroup* K_m *with* $m = \lceil r \rceil + 1$*.*

b) There are finitely many supercuspidal representations σ *of* G *such that* $\text{depth } \sigma \leq r$*.*

Proof a) In the split case this is due to Radhika Ganapathy [12], Lemma 8.2. The proof in the general case is similar to the split one:

Let σ be an irreducible admissible representation of G with $\text{depth } \sigma \leq r$.

Let $\mathcal{C}$ be a chamber of $\mathcal{B}$. For $a \in \Phi$, $a(x)$ is a continuous function on the compact set $\bar{\mathcal{C}}$, and so it has a minimum, which we denote $m(a)$. Define then G_r to be the subgroup of G generated by all the Z_n with $n > r$ and by all U_α with $\alpha = a + \gamma$ satisfying $\gamma > r - m(a)$. We shall show that σ has a vector fixed by $K = G_r$.

For each $x \in \mathcal{B}$ there is $y \in \bar{\mathcal{C}}$ and $g \in G$ such that $x = g \cdot y$. Then $G_{x,r} = gG_{y,r}g^{-1}$. Thus if $\text{depth } \sigma \leq r$, there is $x \in \mathcal{B}$ such that $\sigma^{G_{x,r+}} \neq 0$, and therefore also $\sigma^{G_{y,r+}} \neq 0$.

Take now $\alpha = a + \gamma$ such that $\gamma > r - m(a)$. We have $\alpha(y) = a(y) + \gamma > m(a) + (r - m(a)) = r$, and so we see that the subgroup U_α is contained in $G_{y,r+}$.

Since this is true for any such α and also for all Z_n with $n > r$, we conclude that $G_r \subset G_{y,r+}$. Hence $\sigma^{G_r} \neq 0$.

b) Take the subgroup $K = G_r$ from part a). By Theorem 4.2.1 a) there are only finitely many supercuspidals having a vector fixed by K, and so there are also only finitely many supercuspidals of depth $\leq r$. □

Let $\mathbf{G}$ be a quasisplit classical group over a non-archimedean local field F, i.e., G is split SO_{2n+1}, Sp_{2n} or SO_{2n}, or quasisplit (non-split) even orthogonal corresponding to a quadratic extension E/F. By Arthur's work we know the local Langlands correspondence for G (conditionally on the stabilization of the twisted trace formula) and a slightly weaker form of it when $\mathbf{G}$ is even orthogonal. We know that irreducible admissible representations of G lift to self-dual (irreducible admissible) representations of $GL_N(F)$ for a suitable N.

Let ${}^LG$ be the Langlands dual group of G and W'_F the Weil-Deligne group. For a representation π of G we denote $\phi_\pi : W'_F \to {}^LG$ the corresponding Langlands parameter and Π the lift to $GL_N(F)$ with parameter $\iota \circ \phi_\pi : W'_F \to {}^LG \to {}^LGL_N$ (all this being up to outer conjugacy by O_{2n} in the even orthogonal case). Note that depth $\phi_\pi =$ depth $\iota \circ \phi_\pi$.

Proposition 4.2.3 *Fix* $\mathbf{G}$ *and* F *as above. Then for each* $r \geq 0$ *there exists* $s \geq 0$ *such that:*

a) For all irreducible admissible representations π *of* G *if* depth $\phi_\pi =$ depth $\Pi \leq r$, *then* depth $\pi \leq s$.

b) For all irreducible admissible representations π *of* G *if* depth $\pi \leq r$, *then* depth $\phi_\pi =$ depth $\Pi \leq s$.

Proof a) Assume first that G is not an even orthogonal group, so that we know the full local Langlands correspondence for G.

1) We first prove the statement for supercuspidal representations. Let σ be a supercuspidal representation of G. Let ϕ_σ be its Langlands parameter and σ' the lift to $GL_N(F)$. Assume that depth $\phi_\sigma =$ depth $\sigma' \leq r$.

The representation σ' is not necessarily supercuspidal, but there will be a Levi M of $GL_N(F)$ and a supercuspidal representation τ of M so that σ' is a subquotient of the normalized parabolic induction $\mathrm{Ind}_M^{GL_N(F)}\tau$. By the general properties of depth, we have depth $\tau =$ depth $\sigma' \leq r$.

Now, there are only finitely many standard Levis M (including the one $M = GL_N(F)$). By Proposition 4.2.2, for a fixed M there are only finitely many supercuspidal representations τ of M with depth $\tau \leq r$. And for fixed M and τ, the parabolic induction $\mathrm{Ind}_M^{GL_N(F)}\tau$ has only finitely many possible constituents σ' by [7], Corollary 6.3.7. Let Σ' be the finite set of all triples (M, τ, σ') as above, and let Σ be the finite set of supercuspidal representations of $G(F)$ that lift to some σ' belonging to a triple $(M, \tau, \sigma') \in \Sigma'$. Define s as the maximum of the depths of all representations $\sigma_0 \in \Sigma$.

Thus the representation σ with which we started lies in Σ, and so depth $\sigma \leq s$.

2) Now let σ be a supercuspidal representation of some (standard) Levi subgroup M of $G(F)$.

Then $M = \mathbf{G}_{k_0}(F) \times GL_{k_1}(F) \times \cdots \times GL_{k_l}(F)$, where $\mathbf{G}_{k_0}$ is a group of the same type as $\mathbf{G}$ and smaller rank k_0. Correspondingly we have $\sigma = \sigma_0 \times \sigma_1 \times \cdots \times \sigma_l$, where σ_i are supercuspidal representations of G_{k_0} or $GL_{k_i}(F)$. Let ϕ_i be the Langlands parameter of σ_i, σ_0' the lift of σ_0 to suitable $GL_{k_0'}(F)$, and $M' = GL_{k_0'}(F) \times GL_{k_1}(F) \times \cdots \times GL_{k_l}(F)$. Then $\phi = \phi_0 \oplus \phi_1 \oplus \cdots \oplus \phi_l$ is the Langlands parameter of σ.

We have depth $\sigma = \max(\text{depth } \sigma_i)$ and depth $\phi = \max(\text{depth } \phi_i)$. By the depth preservation theorem for GL, we know that depth $\sigma_i =$ depth ϕ_i for $i > 0$. By the first part of the proof we now get the statement of the proposition in this case.

Since there are finitely many standard Levi subgroups, we can define s to be the maximum of numbers s_M obtained above for individual Levis M.

3) Finally, let π be any representation of $G(F)$ with depth $\phi_\pi =$ depth $\Pi \leq r$. There is a standard Levi M of G and a supercuspidal representation σ of M such that π is a subquotient of the normalized induction $\mathrm{Ind}_M^G\sigma$. By the properties of depth we have depth $\sigma =$ depth π. Let ϕ_σ be the Langlands parameter of σ. We

can also see that depth ϕ_σ = depth ϕ_π. By our assumption, this depth $\leq r$, and so depth σ = depth $\pi \leq s$, as we wanted to show.

4) Now it remains to deal with the even orthogonal case. In this case we know the local Langlands correspondence only up to outer conjugacy by O_{2n}. Given a representation π of G, we know that its Langlands parameter will be one of two possible parameters ϕ_1, ϕ_2. These parameters are conjugate by O_{2n} in ${}^LG$, and so depth ϕ_1 = depth ϕ_2. Now we can use the proof above both for ϕ_1 and ϕ_2 and define s to be the maximum of the two numbers s_1, s_2 that we obtain.

b) The proof is entirely analogous to part a), so we shall omit it. □

For $m \in \mathbb{N}$ denote by K_m the principal congruence subgroup of $GL_N(F)$ of level m, i.e., K_m consists of matrices congruent to the identity I modulo $\wp^m$ (where $\wp$ is the prime ideal of $\mathcal{O}_F$).

Corollary 4.2.4 *Let* **G** *be a quasisplit classical group. For each* $m \in \mathbb{N}$ *there exists a compact open subgroup* $K \subset G$ *such that:*

For each self-dual irreducible admissible representation Π *of* $GL_N(F)$ *which descends to a representation* π *of* G *and satisfies* $\Pi^{K_m} \neq 0$*, one has* $\pi^K \neq 0$.

Proof Since $\Pi^{K_m} \neq 0$, we see that depth $\Pi \leq m$. Thus by Proposition 4.2.3, there is s (depending only on m and not on Π) such that depth $\pi \leq s$. By Proposition 4.2.2 a), there is K (again depending only on s) such that $\pi^K \neq 0$. □

4.3 Local Descent

We can also use the construction of local descent to obtain a sharper estimate on the depths in Proposition 4.2.3 in the case of generic representations.

Theorem 4.3.1 *Let* F *be a p-adic field and* $G = SO_{2n+1}(F)$*. Assume that* p *is sufficiently large.*

a) Let Π *be an (irreducible admissible) representation of* $GL_n(F)$ *that descends to a (generic) representation* π *of* G*. Then* depth $\Pi \geq$ depth π.

b) Let π be an (irreducible admissible) generic representation of G and ϕ_π its Langlands parameter. Then depth $\pi \leq$ depth ϕ_π.

Before proving the theorem, let us review the local descent for the case $\mathbf{G} = SO_{2n+1}$, following [18].

Let τ be a self-dual supercuspidal representation of $GL_{2n}(F)$ that descends to $G = SO_{2n+1}(F)$ (rather than to an even orthogonal group). Denote $H = SO_{4n}(F)$. First view $GL_{2n}(F)$ as the Siegel Levi of a parabolic P in H and take the normalized induction $\rho_\tau = \mathrm{Ind}_P^H \tau |\det|^{1/2}$. Let π_τ be the Langlands quotient of ρ_τ.

The group $H = SO_{4n}(F)$ is the group of isometries of a $4n$-dimensional vector space V preserving a symmetric bilinear form b. Fix maximal isometric subspaces V^+ and V^- of V in duality with respect to b. Also fix a maximal flag $0 \subset V_1^+ \subset V_2^+ \subset \cdots \subset V_n^+ = V^+$ and a corresponding basis $\{e_1, \ldots, e_n\}$ such that $V_i^+ = \mathrm{Span}(e_1, \ldots, e_i)$. Let $\{e_{-1}, \ldots, e_{-n}\}$ be the dual basis of V^- and denote $V_i^- = \mathrm{Span}(e_{-1}, \ldots, e_{-i})$. Finally, let $W = (V_{n-1}^+ + V_{n-1}^-)^\perp$.

Let $P_{n-1} = M_{n-1}N_{n-1}$ be the parabolic subgroup of H stabilizing the flag $0 \subset V_1^+ \subset V_2^+ \subset \cdots \subset V_{n-1}^+$. Then M_{n-1} is isomorphic to $SO(W) \times GL_1(F)^{n-1}$, note that $SO(W) \simeq SO_{2n+2}(F)$.

Fix $\alpha \in F^\times$ and denote $y = e_{2n} - \alpha^2 e_{-2n}$. Let L_{n-1} be the stabilizer of y in $SO(W)$, we have $L_{n-1} = SO(\{y\}^\perp \cap W) \simeq SO_{2n+1}(F) (\subset SO_{2n+2}(F))$ (for more detail on this embedding, see Section 4.5).

Let ψ be an unramified additive character of F and let ψ_{n-1} be the corresponding character of N_{n-1}. For a smooth representation π of H we shall denote by $J_{\psi_{n-1}}(\pi)$ the twisted Jacquet module of π with respect to $N_{n-1}(F)$ and its character ψ_{n-1}, i.e., $J_{\psi_{n-1}}(\pi) = V_\pi / \mathrm{Span}\{\pi(a)v - \psi_{n-1}(a)v | a \in N_{n-1}, v \in V_\pi\}$. Note that, as a representation of the Levi M_{n-1}, this is the same as the (usual, non-twisted) Jacquet module $J(\pi \otimes \psi_{n-1}^{-1}) = V_\pi / \mathrm{Span}\{\pi(a)\psi_{n-1}(a)^{-1}v - v | a \in N_{n-1}, v \in V_\pi\}$.

Finally, let $\sigma = \sigma_{\psi,n-1}(\tau)$ be the restriction of the twisted Jacquet module to L_{n-1}, $\sigma = \mathrm{Res}_{L_{n-1}} J_{\psi_{n-1}}(\pi_\tau)$.

Then σ is the descent of τ and it is an irreducible generic supercuspidal representation of $L_{n-1} \simeq SO_{2n+1}(F)$.

Let us now prove Theorem 4.3.1; note that the local descent construction applies in the cases of other classical groups as well, only generally it may happen that σ is not irreducible, and the descent is in fact an irreducible subquotient of σ. The result and its proof are similar, so we deal only with the case of $G = SO_{2n+1}$ for simplicity of notation.

Proof a) It suffices to prove the theorem only for supercuspidal representations, in general one proceeds as in the proof of Proposition 4.2.3, part 3).

So let us take a supercuspidal self-dual representation τ of $GL_{2n}(F)$ that descends to $G = SO_{2n+1}(F)$. The character $|\det|^{1/2}$ is trivial on $GL_{2n}(\mathcal{O})$, and so it has depth zero, and hence depth $\tau =$ depth $\tau|\det|^{1/2}$. Since parabolic induction preserves depth and π_τ is an irreducible subquotient of $\rho_\tau = \mathrm{Ind}_P^H \tau|\det|^{1/2}$, we see that depth $\tau =$ depth π_τ.

The character ψ_{n-1}^{-1} is unramified, and so depth $\pi_\tau =$ depth $(\pi_\tau \otimes \psi_{n-1}^{-1})$. Since depth is preserved under taking a Jacquet quotient, we see that depth $\tau =$ depth $(\pi_\tau \otimes \psi_{n-1}^{-1}) =$ depth $(J_{\psi_{n-1}}(\pi_\tau))$, where we consider $J = J_{\psi_{n-1}}(\pi_\tau)$ as a representation of M_{n-1}.

We are assuming that the residual characteristic p is large enough, and so the Moy-Prasad filtration on $L_{n-1} \simeq SO_{2n+1}(F)$ is obtained as the intersection of the filtration on $M_{n-1} \simeq SO_{2n+2}(F)$ with L_{n-1} (see [35], 2.3.10.3). Clearly, if J has a vector fixed by some subgroup $(M_{n-1})_{x,r}$, then it has also a vector fixed by $(M_{n-1})_{x,r} \cap L_{n-1}$. Thus depth $\mathrm{Res}_{L_{n-1}} J \leq$ depth $J =$ depth τ, as we wanted to prove.

b) Easily follows from a). □

Let us note that the reason for why we get only one inequality towards depth preservation is the last step of taking the restriction, which may generally decrease the depth. It seems plausible that studying the local descent more closely can prove the full depth preservation for generic representations – we outline a partial progress in this direction in Section 4.5.

4.4 Depth Preservation and Wildly Ramified Groups

In this section we give an example when depth is not preserved under Langlands reciprocity and functoriality. This occurs in some of the examples of residual characteristic $p = 2$ discussed in Section 7.2 of [9]. The fact that depth need not be preserved when residual characteristic is small has already been observed by Gross-Reeder [14]; Reeder-Yu [28] recently provided more examples of this phenomenon.

Let $F = \mathbb{Q}_p$ and $E = \mathbb{Q}_p(\sqrt{p})$. Let $\mathbf{G}$ be the non-split, quasisplit orthogonal group SO^*_{2n} attached to the quadratic extension E/F, let $G = \mathbf{G}(F)$.

Let $\eta = \eta_{E/F} : F^\times \to \mathbb{C}$ be the quadratic character associated to E/F by local class field theory. Note that η is defined by $\ker \eta = N_{E/F}(E^\times)$.

It is easy (and classical) to observe that η is ramified and to compute its depth:

Lemma 4.4.1 *η is ramified. If $p > 2$, then* depth $\eta = 0$. *If $p = 2$, then* depth $\eta = 2$.

Proof We want to check that $\eta|\mathbb{Z}_p^\times \neq 1$. Take $a \in \mathbb{Z}_p^\times$ such that a is not a square modulo p. We shall show that a is not a norm. Assume false, i.e., we have $a = N(x + \sqrt{p}y) = x^2 - py^2$ for some $x, y \in \mathbb{Q}_p$. Let p^m be the common denominator of x and y and write $x = p^{-m}b, y = p^{-m}c$ with $b, c \in \mathbb{Z}_p$ and $m \geq 0$. We obtain $p^{2m}a = b^2 - pc^2$.

If $m > 0$, then p divides b and we can write $b = pd$ and we get $p^{2m-1}a = pd^2 - c^2$. Thus $c = pe$ and $p^{2m-2}a = d^2 - pe^2$. Proceeding in this way, we eventually obtain an equation with $m = 0$.

Let's thus assume that $m = 0$ and $a = b^2 - pc^2$. We see that if a is a norm, it has to be a square modulo p.

Hence η is non-trivial on $\mathbb{Z}_p^\times$, i.e., η is ramified.

The level r subgroup G_r of the Moy-Prasad filtration on $\mathbb{Q}_p^\times$ is just $1+p^{\lceil r \rceil}\mathbb{Z}_p$, where $\lceil r \rceil$ is the smallest integer with $\lceil r \rceil \geq r$. We then have $G_{r+} = \bigcup_{s>r} G_s = 1 + p^{r+1}\mathbb{Z}_p$ for $r \in \mathbb{N}$, and $G_{r+} = G_r = 1 + p^{\lceil r \rceil}\mathbb{Z}_p$ for $r \notin \mathbb{N}$.

When $p > 2$, we can see by Hensel's Lemma that each element of $1 + p\mathbb{Z}_p$ is a norm, and so depth $\eta = 0$. When $p = 2$, every element of $1 + 8\mathbb{Z}_2$ is a norm, whereas 5 is not a norm. Hence depth $\eta = 2$. □

Let now $p = 2$. Take any irreducible, admissible and generic representation π of G of depth zero and let Π be its lift to $GL_{2n}(\mathbb{Q}_2)$. The central character of Π is η, and so depth $\Pi \geq$ depth $\eta = 2$. In fact, in the explicit cases discussed in Section 7.2 of [9], we see that depth $\Pi = 2$ (take for example $\mu_1 = \cdots = \mu_n = 1$, then Π is just the induction of the depth one character $(1, \ldots, 1, \eta, 1, \ldots, 1)$ with η at the n-th coordinate, of the torus $GL_1^{2n}(\mathbb{Q}_2)$).

Note that Π is not unramified, even when π was. This is no contradiction, as the fact that unramified representations lift to unramified ones (via Satake isomorphism) holds only for quasisplit groups which split over an unramified extension, i.e., an unramified group, which is not our case. To sum it up, we have:

Corollary 4.4.2 *Let π be an irreducible, admissible, generic, unramified representation of the ramified group $G = SO_{2n}^*(\mathbb{Q}_2)$ and Π its functorial lift to $GL_{2n}(\mathbb{Q}_2)$. Then Π is not unramified and* depth $\pi = 0 < 2 \leq$ depth Π.

4.5 Restriction and Depth

In the local descent construction in the previous section, we obtained a representation $J_{\psi_{n-1,\alpha}}(\pi_\tau)$ of $M_{n-1} \simeq SO_{2n+2}(F) \times (GL_1(F))^{n-1}$ and restricted it to $SO_{2n+1}(F) \subset SO_{2n+2}(F)$. The depth of the representation may have decreased in this step. The representation of M_{n-1} is a product $\sigma \otimes \chi_1 \otimes \cdots \otimes \chi_{n-1}$, where σ is a representation of $SO_{2n+2}(F)$ and χ_i are characters of $GL_1(F)$.

Hence if we wanted to show that depth does not decrease under this restriction, first we would need to know what are the characters χ_i, e.g., whether they are maybe always unramified in the descent to SO_{2n+1}. The author doesn't know this at present.

Second, we need to understand the restriction of σ from $SO_{2n+2}(F)$ to $SO_{2n+1}(F)$. In this section we show that the depth indeed does not change under this restriction. Let us discuss now the situation slightly more generally:

In the rest of this section let F be a p-adic field, $p \neq 2$. We will consider the (split) orthogonal group of a quadratic space V. Let V be an N-dimensional F-vector space. If $N = 2n$, choose a basis $x_1, \ldots, x_n, x_{-1}, \ldots, x_{-n}$ of V and equip V with the quadratic form $Q(\sum_{i=-n}^{n} a_i x_i) = \sum_{i=1}^{n} a_i a_{-i}$. If $N = 2n+1$, choose a basis $x_0, x_1, \ldots, x_n, x_{-1}, \ldots, x_{-n}$ of V and equip V with the quadratic form $Q(\sum_{i=-n}^{n} a_i x_i)$ $= a_0^2 + \sum_{i=1}^{n} a_i a_{-i}$. In both cases $SO(V) = \{g \in SL(V) | gv = v \text{ for all } v \in V\} = SO_N(F)$.

We'll be considering the embeddings $SO_{2n-1}(F) \subset SO_{2n}(F) \subset SO_{2n+1}(F)$ defined as follows: Take a vector $v \in V$ such that $Q(v) \neq 0$ and take $W = \{v\}^{\perp}$. Then $SO(W) = SO_{N-1}(F)$ is the stabilizer of v in $SO(V)$.

From now on assume that $N = 2n$ is even and take $v = x_n - a^2 x_{-n}$ for some $a \in F^{\times}$ (this is the situation that occurs in the local descent from GL_{2n-2} to SO_{2n-1}).

Proposition 4.5.1 *Let $SO(W) = SO_{2n-1}(F) \subset SO(V) = SO_{2n}(F)$ be an embedding as above. Then there is a quadratic space $U \supset V$ with a quadratic form extending the one on V (hence we will also denote it by Q) such that $SO(W) \times GL_1(F)$ is a Levi subgroup of $SO(U) = SO_{2n+1}(F)$ (where we take $SO(W) \subset SO(U)$ by the natural inclusion). Moreover, $GL_1(F) \cap SO(V) = \{1\}$.*

Proof Take U to be the vector space with basis $x_0, x_1, \ldots, x_n, x_{-1}, \ldots, x_{-n}$ (where $x_1, \ldots, x_n, x_{-1}, \ldots, x_{-n}$ is a basis of V and $x_0 \notin V$), extend the quadratic form to U by $Q(\sum_{i=0}^{2n} a_i x_i) = a_0^2 + \sum_{i=1}^{n} a_i a_{n+i}$. Then V is the orthogonal complement of x_0 in U, which gives $SO(V) = SO_{2n}(F) \subset SO(U) = SO_{2n+1}(F)$.

To show that $SO(W) \times GL_1(F)$ is a Levi in $SO(U) = SO_{2n+1}(F)$, we want to find a corresponding isotropic flag $0 \subset U_0 \subset U$ fixed by a parabolic P. Let U_0 be the 1-dimensional subspace spanned by $u_0 = ax_0 + x_n - a^2 x_{-n}$. We have $Q(u_0) = a^2 - a^2 = 0$,

and so U_0 is isotropic and $SO(U_0) = GL_1(F)$. By definition, $SO(W)$ fixes x_0 and $x_n - a^2 x_{-n}$, and so $SO(W) \times GL_1(F) \subset P$ is the Levi. $\square$

Proposition 4.5.2 *Let π be an irreducible admissible representation of $G = SO_{2n}(F) = SO(V)$ such that its restriction π' to $H = SO(W) = SO_{2n-1}(F) \subset G$ is also irreducible (and automatically admissible). Assume moreover that π' is supercuspidal and generic.*

a) Let χ be a unitary character of $F^\times$ such that $\chi^2 \neq \mathbf{1}$. Then $\mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi \simeq \pi \otimes \chi$.

b) Let K be a compact open subgroup of G. Then $\pi^K \neq \{0\}$ if and only if $\pi'^{K \cap H} \neq \{0\}$.

Proof a) Take U as in the previous proposition so that $H \times GL_1(F)$ is a Levi subgroup of $SO(U) = SO_{2n+1}(F)$, where we take $H = SO(W) \subset G = SO(V) \subset SO(U)$ by the natural inclusions. Consider the parabolic induction $\mathrm{Ind}_{H \times GL_1(F)}^{SO(U)} \pi' \otimes \chi$. By Lemma 2.4.2, this induced representation is irreducible.

Now

$$\mathrm{Ind}_{H \times GL_1(F)}^{SO(U)} \pi' \otimes \chi = \mathrm{Ind}_{G \times GL_1(F)}^{SO(U)} \mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi$$

is irreducible, and so $\mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi$ is also irreducible.

By Frobenius reciprocity we get that

$$\mathrm{Hom}_{G \times GL_1(F)}(\pi \otimes \chi, \mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi) \simeq \mathrm{Hom}_{H \times GL_1(F)}(\pi' \otimes \chi, \pi' \otimes \chi),$$

because the restriction of $\pi \otimes \chi$ to $H \times GL_1(F)$ is $\pi' \otimes \chi$. Since $\pi' \otimes \chi$, $\pi \otimes \chi$, and $\mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi$ are all irreducible, we see that $\mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi \simeq \pi \otimes \chi$.

b) If $\pi^K \neq \{0\}$, then take a non-zero vector v fixed by K. Then clearly $v \in \pi'^{K \cap H}$.

Assume now that $\pi'^{K \cap H} \neq \{0\}$. Let K_0 be an open compact subgroup of $F^\times$ such that $\chi(K_0) = 1$. Then using part a) we have

$$\{0\} \neq \mathrm{Ind}_{(K \cap H) \times K_0}^{K \times K_0} \pi'^{K \cap H} \otimes \chi^{K_0} = \left(\mathrm{Ind}_{H \times GL_1(F)}^{G \times GL_1(F)} \pi' \otimes \chi \right)^{K \times K_0} = (\pi \otimes \chi)^{K \times K_0}.$$

Since $\chi(K_0) = 1$, the space $(\pi \otimes \chi)^{K \times K_0}$ is equal to π^K, and so π^K is also non-zero. $\square$

5. Upper Bound

In this chapter we prove an upper bound on the number $N_{\mathrm{sd}}^K(\lambda)$ of self-dual cuspforms on $GL_N(\mathbb{A}_{\mathbb{Q}})$. The idea of the proof is to consider the functorial descent of each self-dual Π to an appropriate classical group. First we collect some information about functoriality in Section 5.1. The proof is then in Section 5.2.

Throughout this chapter, all groups and representations are adelic, unless stated otherwise.

5.1 Tools

Throughout this chapter, let $K = K_\infty \times \prod K_p$ be a compact subgroup of $GL_N(\mathbb{A})$ with $K_\infty = O(N)$. Mostly we will be working with the principal congruence subgroup $K = K(M)$, for which K_p is the group of matrices $A \in GL_N(\mathcal{O}_p)$ satisfying $A \equiv I \pmod{p^{m(p)}}$ (when $m(p) = 0$ this just means that $K_p = GL_N(\mathcal{O}_p)$), where $M = \prod p^{m(p)}$.

Let us first establish global analogues of the local results from Chapter 4.

Proposition 5.1.1 *a) For every K there is a constant C_K (depending only on K) such that for every automorphic representation Π of $GL_N(\mathbb{A})$ one has $\dim \Pi^K \leq C_K$.*

b) Let $\mathbf{G}$ be a quasisplit classical group. For each $M \in \mathbb{N}$ there exists an arithmetic compact subgroup $K' \subset G(\mathbb{A})$ such that:

For each self-dual cuspidal automorphic representation Π of $GL_N(\mathbb{A})$ which descends to a cuspidal automorphic representation π of $G(\mathbb{A})$ and satisfies $\Pi^{K(M)} \neq 0$, one has $\pi^{K'} \neq 0$.

Proof a) Take Π such that $\Pi^K \neq 0$ and write it as a restricted tensor product $\Pi = \otimes \Pi_v$. Then $\Pi^K = \prod \Pi_v^{K_v}$, and so $\dim \Pi^K = \prod \dim \Pi_v^{K_v}$. When K_v is maximal

and Π_v is unramified (or when $v = \infty$), then $\dim \Pi_v^{K_v} = 1$, and so this is just a finite product.

At each of the remaining finitely many places $v = p$, we have a constant $c_p = c_{p,K_p}$ (independent of Π_p) from Theorem 4.2.1 such that $\dim \Pi_p^{K_p} \leq c_p$. Let C_K be the product of c_p over these ramified places. Then $\dim \Pi^K \leq C_K$ and C_K depends only on K.

b) Take $\Pi = \otimes \Pi_v$ such that $\Pi^K \neq 0$ and which descends to a cuspidal automorphic representation π of $\mathbf{G}(\mathbb{A}_{\mathbb{Q}})$. When K_p is maximal, Π_p is unramified. Thus also π_p is unramified and we can set K_p' to be a special maximal compact subgroup adopted to a standard minimal parabolic so that we have the corresponding Iwasawa decomposition (when $\mathbf{G}$ is split, we can take $K_p' = \mathbf{G}(\mathcal{O}_p)$). Likewise, we can define K_∞' to be a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ (again chosen suitable to provide Iwasawa decomposition).

At the remaining places $v = p$, we use Corollary 4.2.4, which gives us a compact open subgroup $K_p' \subset G$ with $\pi_p^{K_p'} \neq 0$. Then $K' = K_\infty' \times \prod K_p'$ satisfies the conditions of the proposition. □

Let us next discuss to which classical groups $\mathbf{G}$ can Π descend. When $N = 2n+1$ is odd, the only possibility is $\mathbf{G} = Sp_{2n}$. But when $N = 2n$ is even, $\mathbf{G}$ can be split SO_{2n} or SO_{2n+1}, or one of the infinitely many non-split quasisplit groups SO_{2n}^*. However, for a fixed compact subgroup K, only finitely many of these can occur:

Lemma 5.1.2 *Let $K = K(M)$. There is a finite set S_K of quasisplit classical groups $\mathbf{G}$ such that every self-dual cuspidal automorphic representation Π of $GL_N(\mathbb{A})$ with $\Pi^K \neq \{0\}$ descends to some $\mathbf{G} \in S_K$.*

Proof We just need to consider the case when $N = 2n$. Let Π be a self-dual cuspidal automorphic representation of $GL_N(\mathbb{A})$ with $\Pi^K \neq \{0\}$ and let χ be the central character of Π. This is a character of the center $Z \simeq GL_1(\mathbb{A})$ of $GL_N(\mathbb{A})$. Assume moreover that Π descends to one of the non-split groups SO_{2n}^*. This group is then uniquely determined by the central character χ.

We have $\chi(Z \cap K) = 1$, which implies that the conductor of χ divides M. By class field theory, there are only finitely many characters with conductor dividing M. Thus there are also only finitely many possibilities for the group SO^*_{2n}. $\square$

5.2 Statement and Proof of Theorem 5.2.1

We are now ready to prove the upper bound.

For $\lambda > 0$ we have defined $N^K(\lambda) = N^K_{GL(N)}(\lambda) = \sum_{\lambda(\Pi)\leq\lambda} \dim \Pi^K$, where the sum is over all cuspidal automorphic representations Π of $GL_N(\mathbb{A})$ such that the Laplacian eigenvalue $\lambda(\Pi)$ is at most λ. Similarly define $N^K_{\mathrm{sd}}(\lambda)$ to be a similar sum $\sum_{\lambda(\Pi)\leq\lambda} \dim \Pi^K$, but this time ranging only over self-dual representations Π.

Theorem 5.2.1 *Let $K = K(M)$. There are positive constants c and λ_0 (both depending on K) such that $N^K_{\mathrm{sd}}(\lambda) \leq c\lambda^{d/2}$ for all $\lambda > \lambda_0$. Here $d = n^2 + n$ for $N = 2n + \varepsilon$ with $\varepsilon = 0, 1$.*

Proof First of all, when $N = 2$, we can just use the upper bound from (full) Weyl's Law for GL_2, since the dimension d is 2 in both cases. Assume from now on that $N \neq 2$.

We need to estimate

$$N^K_{\mathrm{sd}}(\lambda) = \sum_{\lambda(\Pi)\leq\lambda,\ \Pi\ \mathrm{sd}} \dim \Pi^K.$$

By Lemma 5.1.2, there is a finite set $S = S_K$ such that each of our self-dual Π descends to some $\mathbf{G} \in S$. Hence we can write $N^K_{\mathrm{sd}}(\lambda) = \sum_{\mathbf{G}\in S} N^K_{\mathrm{sd},\mathbf{G}}(\lambda)$, where $N^K_{\mathrm{sd},\mathbf{G}}(\lambda)$ is the sum over all Π which descend to $\mathbf{G}$. It thus suffices to obtain an upper bound for each $N^K_{\mathrm{sd},\mathbf{G}}(\lambda)$.

By Proposition 5.1.1a), there is c_1 such that $\dim \Pi^K \leq c_1$ for each such Π. Hence

$$N^K_{\mathrm{sd},\mathbf{G}}(\lambda) \leq c_1 \cdot \#\{\Pi \mid \lambda(\Pi) \leq \lambda,\ \Pi^K \neq 0,\ \Pi \text{ descends to } \mathbf{G}\}.$$

Let Π be such that $\lambda(\Pi) \leq \lambda$, $\Pi^K \neq 0$, and Π descends to $\mathbf{G}$, and let π be its descent, which is a cuspidal automorphic representation of $\mathbf{G}(\mathbb{A})$. By Theorem 3.3.2,

the Laplacian eigenvalue of Π is $\lambda(\Pi) = c_G\lambda(\pi) + d_G$. Hence there is c_2 such that for sufficiently large λ we have $\lambda(\pi) \leq c_2\lambda$. By Proposition 5.1.1b), there is a compact subgroup K' of $\mathbf{G}(\mathbb{A})$ such that $\pi^{K'} \neq 0$. (Note that neither c_2 nor K' depend on Π.)

Thus we see that

$$\#\{\Pi \mid \lambda(\Pi) \leq \lambda,\ \Pi^K \neq 0, \Pi \text{ descends to } \mathbf{G}\} \leq$$

$$\leq \#\{\pi \mid \pi \text{ representation of } \mathbf{G}(\mathbb{A}),\ \lambda(\pi) \leq c_2\lambda,\ \pi^{K'} \neq 0\}.$$

This is at most $N_{\mathbf{G}}^{K'}(c_2\lambda)$.

Since $N \neq 2$, $\mathbf{G}$ is not SO_2^*. Hence $\mathbf{G}^1(\mathbb{R})$ is not compact and we can use Donnelly's upper bound from Proposition 2.3.4. It gives us $N_{\mathbf{G}}^{K'}(c_2\lambda) \leq c_3\lambda^{d(G)/2}$ for λ large enough and some c_3 as we needed. Here $d(G)$ is the dimension of the symmetric space for $\mathbf{G}$, given by the table preceding Proposition 2.3.4. $\square$

6. Lower Bound

In this chapter we prove a lower bound on the number $N_{\mathrm{sd}}^K(\lambda)$ of self-dual cuspforms on $GL_N(\mathbb{A}_{\mathbb{Q}})$. The idea of the proof is similar to that of the upper bound (i.e., obtaining the self-dual representations as functorial lifts from classical groups), but we have to overcome two additional difficulties: first, a cuspidal automorphic representation on a classical group need not lift to a cuspidal representation on GL_N. We deal with this issue in Section 6.1. And second, we need to obtain an estimate on the size of (K-fixed elements of) global packets. However, to obtain a lower bound it suffices to consider only the case when K is a maximal compact subgroup at all places, which greatly simplifies dealing with the corresponding packets – the result that we need in this case was already discussed in Lemma 2.2.5. The proof of the lower bound is then in Section 6.2.

Let us note that in this chapter we need to use Weak Weyl's Law (formulated in Theorem 2.3.2) to obtain a lower bound on the number $N_G^K(\lambda)$ of cusp forms on a classical group $\mathbf{G}$. This seems not to have appeared in literature yet, but it seems to follow from other results towards Weyl's Law (see the remarks immediately preceding and following Theorem 2.3.2).

Throughout this chapter, all groups and representations are adelic, unless stated otherwise.

6.1 Non-Cuspidal Lifts

Let $\mathbf{G} = \mathbf{G}_{N'}$ be a split group $SO_{N'}$ or $Sp_{N'}$ for $N' = 2n+1$ or $N' = 2n$, respectively (note that we are not considering the even special orthogonal groups). Denote by N the size of the corresponding dual group, i.e., $N = N' - 1 = 2n$ if $N' = 2n+1$ is odd and $N = N' + 1 = 2n+1$ if $N' = 2n$ is even.

Take $\lambda > 0$ and the compact subgroup $K = K_\infty \times \prod K_p$ of $\mathbf{G}(\mathbb{A})$ with $K_p = \mathbf{G}(\mathcal{O}_p)$ and K_∞ a fixed maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ (adopted to a standard Borel subgroup so that we have the corresponding Iwasawa decomposition). Denote by $N^K_{G,\ \mathrm{nc}}(\lambda)$ the sum $\sum_{\lambda(\pi)\leq\lambda} \dim \pi^K$, ranging only over cuspidal automorphic representations π such that their lift Π to $GL_N(\mathbb{A})$ is not cuspidal.

It follows from Theorem 2.2.4 that such a representation Π is an isobaric sum $\Pi = \Pi_1 \boxplus \cdots \boxplus \Pi_k$, where $\sum N_i = N$ and each Π_i is an automorphic representation in the discrete spectrum of $GL_{N_i}(\mathbb{A})$.

Moreover, all the Π_i are self-dual and are lifts of automorphic representations π_i in the discrete spectrum of $\mathbf{G}_{N_i'}(\mathbb{A})$. Here again $N_i = N_i' \pm 1$ depending on the parity of N_i' as at the beginning of this section. Note that all the N_i' have the same parity as N'. By conjugating if necessary, we can assume that the isobaric sum is with respect to the standard Levi $M = \prod_i GL_{N_i}(\mathbb{A})$.

We shall use these facts to estimate the number $N^K_{G,\ \mathrm{nc}}(\lambda)$. For that we again need to relate Laplacian eigenvalues of π and π_i.

Lemma 6.1.1 *Let $M = \prod_i GL_{N_i}(\mathbb{A})$ be a standard Levi subgroup of $GL_N(\mathbb{A})$.*

a) There are constants $c_i > 0$ and d (independent of π) so that $\lambda(\pi) = \sum_i c_i \lambda(\pi_i) + d$ for every cuspidal automorphic representation π of $\mathbf{G}_{N'}(\mathbb{A})$ which lifts to an isobaric sum $\Pi = \Pi_1 \boxplus \cdots \boxplus \Pi_k$ with respect to the Levi M. Here π_i is a representation in the discrete spectrum of $\mathbf{G}_{N_i'}(\mathbb{A})$ which lifts to Π_i.

b) There are constants $c > 0$ and e such that $\lambda(\pi_i) \leq c\lambda(\pi) + e$ for all such π and for all i.

Proof The proof of part a) is similar to the proof of Theorem 3.3.2: Let φ and φ_i be the Langlands parameters of π_∞ and $\pi_{i,\infty}$. Let ι be the embedding $G^\vee \hookrightarrow GL$. Here we abuse notation a little by not distinguishing between the embeddings ι for groups of different size. Then $\iota \circ \varphi$ and $\iota \circ \varphi_i$ are Langlands parameters of Π_∞ and $\Pi_{i,\infty}$. By the properties of an isobaric sum, we have $\iota \circ \varphi = \oplus_i\ \iota \circ \varphi_i$.

Let μ and μ_i be infinitesimal characters of π_∞ and $\pi_{i,\infty}$. Then $\iota(\mu)$ and $\iota(\mu_i)$ are infinitesimal characters of Π_∞ and $\Pi_{i,\infty}$. We see that $\iota(\mu) = \oplus_i\ \iota(\mu_i)$. Evaluating

this infinitesimal character on the Casimir element as in the proof of Theorem 3.3.2 proves our claim.

Part b) now easily follows from a) using the fact that each $\lambda(\pi_i) \geq 0$. $\square$

We are now ready to prove the following proposition:

Proposition 6.1.2 *Let $K = K_\infty \times \prod K_p$ be a compact subgroup of $\mathbf{G}_{N'}(\mathbb{A})$ with $K_p = \mathbf{G}_{N'}(\mathcal{O}_p)$ and K_∞ a fixed maximal compact subgroup of $\mathbf{G}_{N'}(\mathbb{R})$ (adopted to a standard Borel subgroup so that we have the corresponding Iwasawa decomposition). There are positive constants c and λ_0 such that $N^K_{G,\ \mathrm{nc}}(\lambda) \leq c\lambda^{d_0/2}$ for all $\lambda > \lambda_0$.*

Here

- $d_0 = c = 0$ *if* $N' = 1, 3$,
- $d_0 = n^2 - n$ *if* $N' = 2n \geq 2$,
- $d_0 = n^2 - n + 2$ *if* $N' = 2n + 1 \geq 5$.

Note that in all of these cases, d_0 is strictly smaller than the dimension d from Weak Weyl's Law for $\mathbf{G}$.

Proof Denote by $N^K_{G,\ \mathrm{nc},\ M}(\lambda)$ the sum $\sum_{\lambda(\pi)\leq\lambda} \dim \pi^K$, ranging only over cuspidal automorphic representations π which lift to isobaric sums $\Pi = \Pi_1 \boxplus \cdots \boxplus \Pi_k$ with respect to the Levi M. We see that $N^K_{G,\ \mathrm{nc}}(\lambda)$ is a finite sum $\sum_M N^K_{G,\ \mathrm{nc},\ M}(\lambda)$, the sum ranging over standard Levis M. Thus it suffices to estimate $N^K_{G,\ \mathrm{nc},\ M}(\lambda)$.

Since K is maximal at every place, $\dim \pi^K \leq 1$ for every π. Thus

$$N^K_{G,\ \mathrm{nc},\ M}(\lambda) = \#\{\pi \mid \lambda(\pi) \leq \lambda,\ \pi^K \neq 0,$$

$$\pi \text{ lifts to } \Pi_1 \boxplus \cdots \boxplus \Pi_k \text{ with respect to } M\}.$$

For such a π, let $\Pi = \Pi_1 \boxplus \cdots \boxplus \Pi_k$ be its functorial lift and let π_i be the automorphic representations in the discrete spectrum of $G_{N_i'}$ which lift to Π_i (since each packet contains at most one K-fixed element by Lemma 2.2.5, π_i are uniquely

determined by π). Note that π_i is spherical at all places (with respect to suitable maximal compact subgroups).

By Proposition 6.1.1 we have $\lambda(\pi_i) \leq c\lambda(\pi) + e$, where c and e are constants independent of π. Hence we see that $N^K_{G,\ \mathrm{nc},\ M}(\lambda)$ is less than or equal to the product $\prod_i N^K_{G(N_i'),\ \mathrm{disc}}(c\lambda + e)$, where $N^K_{G(N_i'),\ \mathrm{disc}}(c\lambda + e)$ is the corresponding sum over the discrete spectrum of $G(N_i')$.

The same upper bound holds for the discrete and cuspidal spectrum, and so we conclude that

$$N^K_{G,\ \mathrm{nc},\ M}(\lambda) \leq c_M \lambda^{\sum d_i/2},$$

where c_M is a constant depending only on M and d_i is the dimension of the symmetric space for $G_{N_i'}$.

When N is even, $\sum d_i$ will be the largest when $k = 2$ and $N_1 = 2$, $N_2 = N - 2$. When N is odd, $\sum d_i$ will be the largest when $k = 3$ and $N_1 = N_2 = 1$, $N_3 = N - 2$. The proposition then follows from Table 2.2. □

We would expect to have an analogue of Proposition 6.1.2 for general K, not only for maximal one. The obstacle is our (at least the author's) present lack of control over the behaviour of K-spherical members of A-packets for general K.

6.2 Statement and Proof of Theorem 6.2.1

Let us now prove the lower bound on the number of self-dual representations.

Theorem 6.2.1 *Let K be a principal congruence subgroup $K(M)$ of $GL_N(\mathbb{A})$. There are positive constants c and λ_0 such that $N^K_{\mathrm{sd}}(\lambda) \geq c\lambda^{d/2}$ for all $\lambda > \lambda_0$. Here $d = n^2 + n$ for $N = 2n + \varepsilon$ with $\varepsilon = 0, 1$.*

Proof Since the dimension d depends only on N and not on K, it suffices to prove the theorem for $K = K_\infty \times \prod K_p$ with $K_p = GL_N(\mathcal{O}_p)$ and $K_\infty = O(N)$. We shall obtain the lower bound by considering lifts only from $\mathbf{G} = \mathbf{G}_{N'} = SO_{2n+1}$ or Sp_{2n}, when $N = 2n$ or $2n + 1$, respectively.

Denote by $K' = K'_\infty \times \prod K'_p$ a compact subgroup of $\mathbf{G}(\mathbb{A})$ with $K'_p = \mathbf{G}(\mathcal{O}_p)$ and K'_∞ a fixed maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ (adopted to a standard Borel subgroup so that we have the corresponding Iwasawa decomposition). By Weak Weyl's Law (Theorem 2.3.2), we have (for λ sufficiently large) $N_G^{K'}(\lambda) \geq c_1 \lambda^{d/2}$ for a positive constant c_1 and $d = n^2 + n$ as above.

Let $N_{G,\,\mathrm{cusp}}^{K'}(\lambda)$ be the sum $\sum_{\lambda(\pi)\leq\lambda} \dim \pi^{K'}$, ranging only over cuspidal automorphic representations π such that their lift Π to $GL_N(\mathbb{A})$ is cuspidal. Combining Proposition 6.1.2 with Weak Weyl's Law, we obtain $N_{G,\,\mathrm{cusp}}^{K'}(\lambda) \geq c_2 \lambda^{d/2}$ for a positive constant c_2.

Since the dimension of the space of K'-fixed vectors of π is 0 or 1, we see that $N_{G,\,\mathrm{cusp}}^{K'}(\lambda)$ is equal to the number of such K'-spherical representations π. Each of these representations lifts to a K-spherical cuspidal automorphic representation Π with Laplacian eigenvalue $\lambda(\Pi) = c_G \lambda(\pi) + d_G$ by Theorem 3.3.2 (for certain constants c_G, d_G independent of π).

As we observed in Lemma 2.2.5, a global A-packet has at most one K'-spherical element, and so the functoriality mapping $\pi \mapsto \Pi$ is an injection. We conclude that

$$N_{\mathrm{sd}}^K(c_G\lambda + d_G) \geq N_{G,\,\mathrm{cusp}}^{K'}(\lambda) \geq c_2 \lambda^{d/2},$$

which proves the theorem. □

Together with Theorem 5.2.1, this finishes the proof of our main Theorem 1.0.1.

Note that when Weyl's Law is known for G (which is the case when $N = 2n$ and $\mathbf{G} = SO_{2n+1}$ by [23]) and K is a maximal compact subgroup, all of the estimates in the proofs of Theorems 5.2.1 and 6.2.1 are in fact asymptotically equalities. Hence we obtain Weyl's Law (without an error term) for self-dual representations of $GL_{2n}(\mathbb{A})$ in this case as:

Corollary 6.2.2 *Let $N = 2n$ be even and $K = K_\infty \times \prod K_p$ with $K_p = GL_N(\mathcal{O}_p)$ and $K_\infty = O(N)$. Then $N_{\mathrm{sd}}^K(\lambda) = c\lambda^{d/2} + o(\lambda^{d/2})$ for an explicit positive constant c and $d = n^2 + n$.*

Proof We just need to check all the estimates in the proofs of upper and lower bounds.

Let us first consider the case $N = 2$. Let Π be an automorphic representation of $GL_2(\mathbb{A})$ with central character ω. Its dual is then $\widetilde{\Pi} \simeq \Pi \otimes \omega^{-1}$. If Π is K-spherical (for maximal K as above), the central character ω is trivial on $\mathcal{O}_p^\times$ for every p. Hence ω is an unramified character of $\mathbb{A}^\times$, and so it is trivial (class field theory attaches to ω an extension of $\mathbb{Q}$ which is unramified everywhere, and so it is just the trivial extension). We see that every K-spherical automorphic representation of $GL_2(\mathbb{A})$ is self-dual, and so our claim is just Weyl's Law in this case.

Take now $N \geq 4$. With notation as in the proof of Theorem 5.2.1, we have $N_{\mathrm{sd}}^K(\lambda) = \sum_{\mathbf{G} \in S} N_{\mathrm{sd},\mathbf{G}}^K(\lambda)$. As we have seen in the proof of 5.2.1, the main term in the sum on the right hand side is the one corresponding to $\mathbf{G} = SO_{2n+1}$, and so $N_{\mathrm{sd}}^K(\lambda) = N_{\mathrm{sd},\ SO(2n+1)}^K(\lambda) + o(\lambda^{d/2})$. From now on, let $\mathbf{G} = SO_{2n+1}$. When Π is unramified everywhere, $\dim \Pi^K = 1$, and so

$$N_{\mathrm{sd},\mathbf{G}}^K(\lambda) = \#\{\Pi \mid \lambda(\Pi) \leq \lambda,\ \Pi^K \neq 0,\ \Pi \text{ descends to } \mathbf{G}\}.$$

Let $K' = K'_\infty \times \prod K'_p$ be a compact subgroup of $\mathbf{G}(\mathbb{A})$ with $K'_p = \mathbf{G}(\mathcal{O}_p)$ and K'_∞ a fixed maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ (adopted to a standard Borel subgroup so that we have the corresponding Iwasawa decomposition).

Take a K-spherical self-dual representation Π and denote by π its descent to $\mathbf{G}(\mathbb{A})$. Then π is K'-spherical, and by Lemma 2.2.5, π is unique. By Theorem 3.3.2, the Laplacian eigenvalue of π is $\lambda(\Pi) = c_G \lambda(\pi) + d_G$ (for suitable constants c_G, d_G).

Hence, with notation as in the proof of 6.2.1, $N_{\mathrm{sd},\mathbf{G}}^K(c_G\lambda + d_G) = N_{G,\ \mathrm{cusp}}^{K'}(\lambda)$ is the number of cusp forms of $\mathbf{G}(\mathbb{A})$ with Laplacian eigenvalue bounded from above by λ, whose lifts to $GL_N(\mathbb{A})$ are cuspidal. By Proposition 6.1.2 and by Weyl's Law for $\mathbf{G}$, we have

$$N_{G,\ \mathrm{cusp}}^{K'}(\lambda) = N_G^{K'}(\lambda) - N_{G,\ \mathrm{nc}}^{K'}(\lambda) = c_1 \lambda^{d/2} + o(\lambda^{d/2})$$

for an explicit constant c_1 (here $N_{G,\ \mathrm{nc}}^{K'}(\lambda)$ denotes the number of cusp forms whose lift is non-cuspidal).

Putting everything together, we obtain

$$N_{\mathrm{sd}}^{K}(c_G\lambda+d_G) = N_{\mathrm{sd},\mathbf{G}}^{K}(c_G\lambda+d_G)+o(\lambda^{d/2}) = N_{G,\ \mathrm{cusp}}^{K'}(\lambda)+o(\lambda^{d/2}) = c_1\lambda^{d/2}+o(\lambda^{d/2}).$$

This finishes the proof when we set $c = c_1 c_G^{-d/2}$. □

A similar statement for K-spherical self-dual representations of $GL_{2n+1}(\mathbb{A})$ would follow from Weyl's Law for $Sp_{2n}(\mathbb{A})$. However, regardless of the parity of N, we are crucially relying on the assumption that K is maximal everywhere. Otherwise we do not know how to obtain sufficiently precise estimates in several of the steps of the proof, namely, for the choice of a corresponding compact subgroup K', for the dimension of K- and K'-fixed vectors, and for the sizes of global A-packets.

LIST OF REFERENCES

LIST OF REFERENCES

[1] Jeffrey Adams and David A Vogan Jr. The contragredient. *arXiv preprint arXiv:1201.0496*, 2012.

[2] Jeffrey Adler. Self-contragredient supercuspidal representations of GL_n. *Proceedings of the American Mathematical Society*, 125(8):2471–2479, 1997.

[3] James Arthur. *The Endoscopic Classification of Representations Orthogonal and Symplectic Groups*, volume 61. Colloquium Publications, American Mathematical Society, 2013.

[4] Mahdi Asgari and Freydoon Shahidi. Functoriality for general spin groups. *Manuscripta Math., published online*, 2014.

[5] Joseph N Bernshtein. All reductive p-adic groups are of type I. *Funkcional. Anal. i Prilozen*, 8(2):3–6, 1974.

[6] Joseph N Bernshtein and Andrei Zelevinskii. Representations of the group $GL(n, F)$ where F is a non-archimedean local field. *Russian Mathematical Surveys*, 31(3):1–68, 1976.

[7] William Casselman. Introduction to the theory of admissible representations of p-adic reductive groups. *online*, 1995.

[8] James W Cogdell, Henry H Kim, Ilya I Piatetski-Shapiro, and Freydoon Shahidi. Functoriality for the classical groups. *Publ. Math. Inst. Hautes Études Sci*, 99:163–233, 2004.

[9] James W Cogdell, Ilya I Piatetski-Shapiro, and Freydoon Shahidi. Functoriality for the quasisplit classical groups. In *On certain L-functions, Clay Math. Proc., Amer. Math. Soc.*, number 13, pages 455–499, 2011.

[10] Harold Donnelly. On the cuspidal spectrum for finite volume symmetric spaces. *J. Differential Geom*, 17(2):239–253, 1982.

[11] William Fulton and Joe Harris. *Representation theory: a first course*, volume 129. Graduate Texts in Mathematics, Springer, 1991.

[12] Radhika Ganapathy. The local Langlands correspondence for GSp_4 over local function fields. *arXiv preprint arXiv:1305.6088*, 2013.

[13] David Ginzburg, Stephen Rallis, and David Soudry. *The descent map from automorphic representations of GL(n) to classical groups*. World Scientific, 2011.

[14] Benedict H Gross and Mark Reeder. Arithmetic invariants of discrete Langlands parameters. *Duke Mathematical Journal*, 154(3):431–508, 2010.

[15] Michael Harris and Richard Taylor. *The Geometry and Cohomology of Some Simple Shimura Varieties*. Number 151. Annals of Mathematical Studies, Princeton University Press, 2001.

[16] Guy Henniart. Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p-adique. *Inventiones Mathematicae*, 139(2):439–455, 2000.

[17] Hervé Jacquet, Ilya Piatetski-Shapiro, and Joseph Shalika. Conducteur des représentations du groupe linéaire. *Mathematische Annalen*, 256(2):199–214, 1981.

[18] Dihua Jiang and David Soudry. Appendix: On the local descent from $GL(n)$ to classical groups. *American Journal of Mathematics*, 134(3):767–772, 2012.

[19] Anthony W Knapp. *Lie groups beyond an introduction*, volume 140. Progress in Mathematics, Birkhäuser, 2002.

[20] Jean-Pierre Labesse and Werner Müller. Weak Weyl's law for congruence subgroups. *Asian Journal of Mathematics*, 8(4):733–746, 2004.

[21] Robert P Langlands. On the classification of irreducible representations of real algebraic groups. *Representation theory and harmonic analysis on semisimple Lie groups*, 31:101–170, 1989.

[22] Erez Lapid and Werner Müller. Spectral asymptotics for arithmetic quotients of $SL(n,\mathbb{R})/SO(n)$. *Duke Mathematical Journal*, 149(1):117–155, 2009.

[23] Elon Lindenstrauss and Akshay Venkatesh. Existence and Weyl's law for spherical cusp forms. *Geometric And Functional Analysis*, 17(1):220–251, 2007.

[24] Chung Pang Mok. Endoscopic classification of representations of quasi-split unitary groups. *Memoirs of the American Mathematical Society, to appear*.

[25] Allen Moy and Gopal Prasad. Unrefined minimal K-types for p-adic groups. *Inventiones mathematicae*, 116(1):393–408, 1994.

[26] Allen Moy and Gopal Prasad. Jacquet functors and unrefined minimal K-types. *Commentarii Mathematici Helvetici*, 71(1):98–121, 1996.

[27] Werner Müller. Weyl's Law for the cuspidal spectrum of SL_n. *Annals of Mathematics*, pages 275–333, 2007.

[28] Mark Reeder and Jiu-Kang Yu. Epipelagic representations and invariant theory. *Journal of American Mathematical Society*, 27(2):437–477, 2014.

[29] David Rohrlich. Self-dual artin representations. In *Automorphic Representation and L-functions, Tata Institute of Fundamental Research Studies in Mathematics*, number 22, pages 455–499, 2013.

[30] Atle Selberg. In *Harmonic Analysis, Göttingen Lecture notes, Collected papers, Vol. I*, pages 626–674. Springer Verlag, 1989.

[31] Freydoon Shahidi. On certain L-functions. *American Journal of Mathematics*, 103(2):297–355, 1981.

[32] Freydoon Shahidi. *Eisenstein series and automorphic L-functions*, volume 58. Colloquium Series, American Mathematical Society, 2010.

[33] Freydoon Shahidi. Arthur packets and the Ramanujan conjecture. *Kyoto Journal of Mathematics*, 51(1):1–23, 2011.

[34] Jacques Tits. Reductive groups over local fields. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977)*, volume 1, pages 29–69, 1979.

[35] Jiu-Kang Yu. Bruhat-Tits theory and buildings. In *Ottawa lectures on admissible representations of reductive p-adic groups, Fields Inst. Monogr.*, number 26, pages 53–77, 2009.

VITA

VITA

Vítězslav Kala received Bachelor's (in 2007) and Master's (in 2009) degrees in Mathematics from Charles University in Prague, Czech Republic. He has been a PhD student in the Department of Mathematics at Purdue University from August 2009 to August 2014. During his first three years at Purdue, he has been supported by the International Fulbright Science and Technology Award.